

METHODS FOR GENERATING HYPERMUTABLE MICROBES

5 This application claims the benefit of provisional application S.N.
60/181,929 filed February 11, 2000.

FIELD OF THE INVENTION

10 The invention is related to the area of mismatch repair genes. In
particular it is related to the field of *in situ* mutagenesis of single celled
organisms.

BACKGROUND OF THE INVENTION

15 Within the past four years, the genetic cause of the Hereditary
Nonpolyposis Colorectal Cancer Syndrome (HNPCC), also known as
Lynch syndrome II, has been ascertained for the majority of kindred's
affected with the disease (14). The molecular basis of HNPCC involves
genetic instability resulting from defective mismatch repair (MMR).
Several genes have been identified in humans that encode for proteins and
20 appear to participate in the MMR process, including the *mutS* homologs
GTBP, *hMSH2*, and *hMSH3* and the *mutL* homologs *hMLH1*, *hMLH3*,
hPMS1, and *hPMS2* (4,9,11,17,19,22,24,38). Germline mutations in four
of these genes (*hMSH2*, *hMLH1*, *hPMS1*, and *hPMS2*) have been identified
in HNPCC kindred's (13). Though the mutator defect that arises from the
25 MMR deficiency can affect any DNA sequence, microsatellite sequences
are particularly sensitive to MMR abnormalities (14, 25, 27, 29).
Microsatellite instability (MI) is therefore a useful indicator of defective
MMR. In addition to its occurrence in virtually all tumors arising in
HNPCC patients, MI is found in a small fraction of sporadic tumors with
30 distinctive molecular and phenotypic properties (13).

 HNPCC is inherited in an autosomal dominant fashion, so that the
normal cells of affected family members contain one mutant allele of the

09780673-1024204

relevant MMR gene (inherited from an affected parent) and one wild-type allele (inherited from the unaffected parent). During the early stages of tumor development, however, the wild-type allele is inactivated through a somatic mutation, leaving the cell with no functional MMR gene and
5 resulting in a profound defect in MMR activity. Because a somatic mutation in addition to a germ-line mutation is required to generate defective MMR in the tumor cells, this mechanism is generally referred to as one involving two hits, analogous to the biallelic inactivation of tumor suppressor genes that initiate other hereditary cancers. In line with this
10 two-hit mechanism, the non-neoplastic cells of HNPCC patients generally retain near normal levels of MMR activity due to the presence of the wild-type allele (11, 13, 24). In addition, similar findings are observed in other diploid organisms (2, 5, 8).

The ability to alter signal transduction pathways by manipulation of
15 a gene product's function, either by over-expression of the wild type protein or a fragment thereof, or by introduction of mutations into specific protein domains of the protein, the so-called dominant-negative inhibitory mutant, were described over a decade ago in the yeast system
Saccharomyces cerevisiae by Herskowitz (Nature 329:219-222, 1987). It
20 has been demonstrated that over-expression of wild type gene products can result in a similar, dominant-negative inhibitory phenotype due most likely to the "saturating-out" of a factor, such as a protein, that is present at low levels and necessary for activity; removal of the protein by binding to a high level of its cognate partner results in the same net effect, leading to
25 inactivation of the protein and the associated signal transduction pathway.

Recently, work done by Nicolaides et.al. (32) has demonstrated the utility of introducing dominant negative inhibitory mismatch repair mutants into mammalian cells to confer global DNA hypermutability. There is a need in the art for additional techniques for generating mutations in
5 bacteria which can be used to make strains for production, biocatalysis, bioremediation, and drug discovery.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide a method for
10 rendering bacterial cells hypermutable.

It is another object of the present invention to provide genetically altered bacteria.

It is yet another object of the present invention to provide a method to produce bacteria that are hypermutable.

15 It is an object of the invention to provide a method to inactivate the process that results in hypermutable cells following strain selection.

It is a further object of the invention to provide a method of mutating a gene of interest in a bacterium.

These and other embodiments of the invention are provided by one
20 or more of the embodiments described below. In one embodiment, a method is provided for making a hypermutable bacteria. A polynucleotide comprising a dominant negative allele of a mismatch repair gene is introduced into a bacterium, whereby the cell becomes hypermutable. Preferably the allele is under the control of an inducible transcription
25 regulatory sequence.

According to another aspect of the invention a homogeneous composition of cultured, hypermutable, bacteria is provided. The bacteria comprise a dominant negative allele of a mismatch repair gene. Preferably the allele is under the control of an inducible transcription regulatory
30 sequence.

Another embodiment of the invention provides a method for generating a mutation in a gene of interest. A bacterial culture comprising the gene of interest and a dominant negative allele of a mismatch repair gene is grown. The cell is hypermutable. It is tested to determine whether the gene of interest harbors a mutation. Preferably the allele is under the control of an inducible transcription regulatory sequence.

According to still another aspect of the invention a method for generating a mutation in a gene of interest is provided. A bacterium comprising the gene of interest and a dominant negative allele of a mismatch repair gene is grown to form a population of mutated bacteria. The population of mutated bacteria is cultivated under trait selection conditions. At least one of the cultivated bacteria is tested to determine that the gene of interest harbors a mutation. Preferably the allele is under the control of an inducible transcription regulatory sequence.

Still another aspect of the invention is a method for enhancing the mutation rate of a bacterium. A bacterium comprising a dominant negative allele of an MMR gene is exposed to a mutagen whereby the mutation rate of the bacterium is enhanced in excess of the rate in the absence of mutagen and in excess of the rate in the absence of the dominant negative allele. Preferably the allele is under the control of an inducible transcription regulatory sequence.

Yet another aspect of the invention is a method for generating an MMR-proficient bacterium with a new output trait. A mismatch repair deficient bacterium comprising a gene of interest and a dominant negative allele of a mismatch repair gene is grown to form a population of mutated bacteria. The population of mutated bacteria is cultivated under trait selection conditions. At least one of the cultivated bacteria is tested to determine that the gene of interest harbors a mutation. Mismatch repair activity is restored to the at least one of the cultivated

bacteria. Preferably the allele is under the control of an inducible transcription regulatory sequence.

These and other embodiments of the invention provide the art with methods that can generate enhanced mutability in bacteria as well as

5 providing prokaryotic organisms harboring potentially useful mutations to generate novel output traits for commercial applications. The ability to create hypermutable organisms using dominant negative alleles has great commercial value for the generation of innovative bacterial strains that display new output features useful for a variety of applications, including

10 but not limited to the manufacturing industry for the generation of new biochemicals useful for detoxifying noxious chemicals from by-products of manufacturing processes or those used as catalysts, as well as helping in remediation of toxins present in the environment, including but not limited to polychlorobenzenes (PCBs), heavy metals and other environmental

15 hazards for which there is a need to remove them from the environment. In addition to obtaining organisms that are useful for removal of toxins from the environment, novel microbes can be selected for enhanced activity to either produce increased quantity or quality of a protein or non-protein therapeutic molecule by means of biotransformation (3).

20 Biotransformation is the enzymatic conversion, by a microbe or an extract derived from the microbe, of one chemical intermediate to the next product. There are many examples of biotransformation in use for the commercial manufacturing of important biological and chemical products, including Penicillin G, Erythromycin, and Clavulanic Acid as well as organisms that

25 are efficient at conversion of "raw" materials to advanced intermediates and/or final products (Berry, A. Trends Biotechnol. 14(7):250-256). The ability to control DNA hypermutability in host bacterial strains using a dominant negative MMR (as described above) allows for the generation of variant subtypes that can be selected for new phenotypes of commercial

30 interest, including but not limited to organisms that are toxin-resistant, have

the capacity to degrade a toxin *in situ* or the ability to convert a molecule from an intermediate to either an advanced intermediate or a final product. Other applications using dominant negative MMR genes to produce genetic alteration of bacterial hosts for new output traits include but are not limited to recombinant production strains that produce higher quantities of a recombinant polypeptide as well as the use of altered endogenous genes that can transform chemical or catalyze manufacturing downstream processes.

This application teaches of the use of a regulatable dominant negative MMR phenotype to produce a prokaryotic strain with a commercially beneficial output trait. Using this process, microbes expressing a dominant negative MMR can be directly selected for the phenotype of interest. Once a selected bacterium with a specified output trait is isolated, the hypermutable activity of the dominant negative MMR allele can be turned-off by several methods well known to those skilled in the art. For example, if the dominant-negative allele is expressed by an inducible promoter system, including but not limited to promoters such as: TAC-LACI, tryp (Brosius et.al. *Gene* 27:161-172, 1984), araBAD (Guzman et.al., *J. Bact.* 177:4121-4130, 1995) pLex (La Vallie et.al., *Bio.Technology* 11:187-193, 1992), pRSET (Schoepfer, R. *Gene* 124:83-85, 1993) , pT7 (Studier *J. Mol. Biol.* 219(1):37-44, 1991) etc., the inducer is removed and the promoter activity is reduced, or a system that excises the MMR gene insert from the host cells harboring the expression vector such as the Cre-lox (Hasan, N. et.al. *Gene* 2:51-56, 1994), as well as methods that can homologously knockout of the expression vector. In addition to the recombinant methods outlined above that have the capacity to eliminate the MMR activity from the microbe, it has been demonstrated that many chemicals have the ability to “cure” microbial cells of plasmids. For example, chemical treatment of cells with drugs including bleomycin (Attfield et al. *Antimicrob. Agents Chemother.* 27:985-988, 1985) or

novobiocin, coumermycin, and quinolones (Fu et al. *Chemotherapy* 34:415-418, 1988) have been shown to result in microbial cells that lack endogenous plasmid as evidenced by Southern analysis of cured cells as well as sensitivity to the appropriate antibiotic (1, 41-43). Whether by use of recombinant means or treatment of cells with chemicals, removal of the MMR-expression plasmid results in the re-establishment of a genetically stable microbial cell-line. Therefore, the restoration of MMR allows host bacteria to function normally to repair DNA. The newly generated mutant bacterial strain that exhibits a novel, selected output trait is now suitable for a wide range of commercial processes or for gene/protein discovery to identify new biomolecules that are involved in generating a particular output trait.

While it has been documented that MMR deficiency can lead to as much as a 1000-fold increase in the endogenous DNA mutation rate of a host, there is no assurance that MMR deficiency alone will be sufficient to alter every gene within the DNA of the host bacterium to create altered biochemicals with new activity(s). Therefore, the use of chemical agents and their respective analogues such as ethidium bromide, EMS, MNNG, MNU, Tamoxifen, 8-Hydroxyguanine, as well as others listed but not limited to in publications by: Khromov-Borisov, N.N., et.al. (*Mutat. Res.* 430:55-74, 1999); Ohe, T., et.al. (*Mutat. Res.* 429:189-199, 1999); Hour, T.C. et.al. (*Food Chem. Toxicol.* 37:569-579, 1999); Hrelia, P., et.al. (*Chem. Biol. Interact.* 118:99-111, 1999); Garganta, F., et.al. (*Environ. Mol. Mutagen.* 33:75-85, 1999); Ukawa-Ishikawa S., et.al. (*Mutat. Res.* 412:99-107, 1998); www.ehs.utah.edu/ohh/mutagens, etc. can be used to further enhance the spectrum of mutations and increase the likelihood of obtaining alterations in one or more genes that can in turn generate host bacteria with a desired new output trait(s) (10, 39, 40). Prior art teaches that mismatch repair deficiency leads to hosts with an increased resistance to toxicity by chemicals with DNA damaging activity. This feature allows

for the creation of additional genetically diverse hosts when mismatch defective bacteria are exposed to such agents, which would be otherwise impossible due to the toxic effects of such chemical mutagens [Colella, G., et.al. (Br. J. Cancer 80:338-343, 1999); Moreland, N.J., et.al. (Cancer Res. 59:2102-2106, 1999); Humbert, O., et.al. (Carcinogenesis 20:205-214, 1999); Glaab, W.E., et.al. (Mutat. Res. 398:197-207, 1998)]. Moreover, prior art teaches that mismatch repair is responsible for repairing chemical-induced DNA adducts, so therefore blocking this process could theoretically increase the number, types, mutation rate and genomic alterations of a bacterial host [Rasmussen, L.J. et.al. (Carcinogenesis 17:2085-2088, 1996); Sledziewska-Gojska, E., et.al. (Mutat. Res. 383:31-37, 1997); and Janion, C. et.al. (Mutat. Res. 210:15-22, 1989)]. In addition to the chemicals listed above, other types of DNA mutagens include ionizing radiation and UV-irradiation, which are known to cause DNA mutagenesis in bacteria can also be used to potentially enhance this process. These agents which are extremely toxic to host cells and therefore result in a decrease in the actual pool size of altered bacterial cells are more tolerated in MMR defective hosts and in turn allow for a enriched spectrum and degree of genomic mutation (7).

This application teaches new uses of MMR deficient bacterial cells to create commercially viable microbes that express novel output traits. Moreover, this application teaches the use of dominant negative MMR genes to decrease the endogenous MMR activity of the host followed by placing the cells under selection to obtain a desired, sought after output trait for commercial applications such as but not limited to recombinant manufacturing, biotransformation and bioremediation. Furthermore, the application teaches the use of restoring MMR activity to the hypermutable bacterial host following strain selection of the variant of interest as a means to genetically "fix" the new mutations in the host genome. The application also teaches the use of enhanced hypermutability in bacteria by using

MMR deficiency and chemical or radiation mutagenesis to create variant subtypes of bacteria useful for commercial and other applications. The application describes uses of hypermutable bacteria for producing strains that can be used to generate new output traits for chemical manufacturing, pharmaceutical and other commercially applicable processes.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Western blot of steady-state PMS134 levels in IPTG-treated samples in DH10B strain. Cells containing the pTACPMS134 (lane 2) showed a robust steady state level of protein after induction in contrast to cells expressing empty vector (lane 1). Blots were probed with an anti-human-PMS2 polyclonal antibody.

Figure 2. Western blot of PMS134V5 and PMSR3V5 in IPTG-treated (+) and untreated (-) samples in BL21 strain. Blots were probed with an anti-V5 antibody, which is directed to the C-terminal tag of each protein.

Figure 3. Number of Kanamycin resistant PMS134 and vector control DH10B clones. IPTG-induced strains were grown and plated onto KAN plates and grown for 18 hours at 37°C to identify number of KAN resistant clones due to genetic alteration.

Figure 4. Number of Kanamycin Resistant PMS134, PMSR3 and vector control BL21 clones. IPTG-induced strains were grown and plated onto AMP and KAN plates and grown for 18 hours at 37°C to identify number of KAN resistant clones due to genetic alteration.

Figure 5. (A) Western blot of steady-state ATPMS134_{flag} in IPTG-treated samples in DH10B. Lysates from untransfected cells (lane 1) and a

bacterial clone expressing the Arabidopsis thaliana PMS134 truncated protein with a FLAG epitope fused to the C-terminus (ATPMS134flag) (lane 2) were electrophoresed on SDS-PAGE gels. Blots were probed with an anti-FLAG monoclonal antibody directed to the FLAG epitope. (B)

5 Number of Kanamycin Resistant ATPMS134flag and vector control DH10B clones. IPTG-induced strains were grown and plated onto AMP and KAN plates and grown for an additional 18 hours at 37°C to identify number of KAN resistant clones due to genetic alteration.

10 **Figure 6.** Generation of high recombinant producer BGAL-MOR lines in PMS134 expressing DH5alpha host strains.

DETAILED DESCRIPTION OF THE INVENTION

15 The inventors present a method for developing hypermutable bacteria by altering the activity of endogenous mismatch repair activity of hosts. Wild type and some dominant negative alleles of mismatch repair genes, when introduced and expressed in bacteria, increase the rate of spontaneous mutations by reducing the effectiveness of the endogenous
20 MMR-mediated DNA repair activity, thereby rendering the bacteria highly susceptible to genetic alterations due to hypermutability. Hypermutable bacteria can then be utilized to screen for novel mutations in a gene or a set of genes that produce variant siblings that exhibit a new output trait(s) not found in the wild type cells.

25 The process of mismatch repair, also called mismatch proofreading, is an evolutionarily highly conserved process that is carried out by protein complexes described in cells as disparate as prokaryotic cells such as bacteria to more complex mammalian cells (14, 29, 31, 33, 34). A mismatch repair gene is a gene that encodes one of the proteins of such a
30 mismatch repair complex. Although not wanting to be bound by any

particular theory of mechanism of action, a mismatch repair complex is believed to detect distortions of the DNA helix resulting from non-complementary pairing of nucleotide bases. The non-complementary base on the newer DNA strand is excised, and the excised base is replaced with the appropriate base that is complementary to the older DNA strand. In this way, cells eliminate many mutations that occur as a result of mistakes in DNA replication, resulting in genetic stability of the sibling cells derived from the parental cell.

Some wild type alleles as well as dominant negative alleles cause a mismatch repair defective phenotype even in the presence of a wild-type allele in the same cell. An example of a dominant negative allele of a mismatch repair gene is the human gene *hPMS2-134*, which carries a truncation mutation at codon 134 (32). The mutation causes the product of this gene to abnormally terminate at the position of the 134th amino acid, resulting in a shortened polypeptide containing the N-terminal 133 amino acids. Such a mutation causes an increase in the rate of mutations, which accumulate in cells after DNA replication. Expression of a dominant negative allele of a mismatch repair gene results in impairment of mismatch repair activity, even in the presence of the wild-type allele. Any mismatch repair allele, which produces such effect, can be used in this invention. In addition, the use of over-expressed wildtype MMR gene alleles from human, mouse, plants, and yeast in bacteria has been shown to cause a dominant negative effect on the bacterial hosts MMR activity (9, 33, 34, 38).

Dominant negative alleles of a mismatch repair gene can be obtained from the cells of humans, animals, yeast, bacteria, plants or other organisms. Screening cells for defective mismatch repair activity can identify such alleles. Mismatch repair genes may be mutant or wild type. Bacterial host MMR may be mutated or not. The term bacteria used in this application include any organism from the prokaryotic *kingdom*. These

organisms include genera such as but not limited to *Agrobacterium*, *Anaerobacter*, *Aquabacterium*, *Azorhizobium*, *Bacillus*, *Bradyrhizobium*, *Cryobacterium*, *Escherichia*, *Enterococcus*, *Heliobacterium*, *Klebsiella*, *Lactobacillus*, *Methanococcus*, *Methanothermobacter*, *Micrococcus*, *Mycobacterium*, *Oceanomonas*, *Pseudomonas*, *Rhizobium*, *Staphylococcus*, *Streptococcus*, *Streptomyces*, *Thermusaquaticus*, *Thermaerobacter*, *Thermobacillus*, etc. Other procaryotes that can be used for this application are listed at (www.bacterio.cict.fr/validgenericnames).

Bacteria exposed to chemical mutagens or radiation exposure can be screened for defective mismatch repair. Genomic DNA, cDNA, or mRNA from any cell encoding a mismatch repair protein can be analyzed for variations from the wild type sequence. Dominant negative alleles of a mismatch repair gene can also be created artificially, for example, by producing variants of the *hPMS2-134* allele or other mismatch repair genes (32). Various techniques of site-directed mutagenesis can be used. The suitability of such alleles, whether natural or artificial, for use in generating hypermutable bacteria can be evaluated by testing the mismatch repair activity (using methods described in ref 32) caused by the allele in the presence of one or more wild-type alleles, to determine if it is a dominant negative allele.

A bacterium that over-expresses a wild type mismatch repair allele or a dominant negative allele of a mismatch repair gene will become hypermutable. This means that the spontaneous mutation rate of such bacteria is elevated compared to bacteria without such alleles. The degree of elevation of the spontaneous mutation rate can be at least 2-fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, or 1000-fold that of the normal bacteria as measured as a function of bacterial doubling/minute.

According to one aspect of the invention, a polynucleotide encoding either a wild type or a dominant negative form of a mismatch repair protein is introduced into bacteria. The gene can be any dominant negative allele

encoding a protein which is part of a mismatch repair complex, for example, *mutS*, *mutL*, *mutH*, or *mutY* homologs of the bacterial, yeast, plant or mammalian genes (14, 28). The dominant negative allele can be naturally occurring or made in the laboratory. The polynucleotide can be in the form of genomic DNA, cDNA, RNA, or a chemically synthesized polynucleotide or polypeptide. The molecule can be introduced into the cell by transfection, transformation, conjugation, fusion, or other methods well described in the literature.

Any process can be used whereby a polynucleotide or polypeptide is introduced into a cell. The process of gene transfer can be carried out in a bacterial culture using a suspension culture. The bacteria can be any type classified under the prokaryotes.

In general, gene transfer will be carried out using a suspension of cells but other methods can also be employed as long as a sufficient fraction of the treated cells incorporate the polynucleotide or polypeptide so as to allow recipient cells to be grown and utilized. The protein product of the polynucleotide may be transiently or stably expressed in the cell. Techniques for gene transfer are well known to those skilled in the art. Available techniques to introduce a polynucleotide or polypeptide into a prokaryote include but are not limited to electroporation, transduction, cell fusion, the use of chemically competent cells (e.g. calcium chloride), and packaging of the polynucleotide together with lipid for fusion with the cells of interest. Once a cell has been transformed with the dominant negative mismatch repair gene or protein, the cell can be propagated and manipulated in either liquid culture or on a solid agar matrix, such as a petri dish. If the transfected cell is stable, the gene will be retained and expressed at a consistent level when the promoter is constitutively active, or when in the presence of appropriate inducer molecules when the promoter is inducible, for many cell generations, and a stable, hypermutable bacterial strain results.

An isolated bacterial cell is a clone obtained from a pool of a bacterial culture by chemically selecting out non-recipient strains using, for example, antibiotic selection of an expression vector. If the bacterial cell is derived from a single cell, it is defined as a clone.

5 A polynucleotide encoding a dominant negative form of a mismatch repair protein can be introduced into the genome of a bacterium or propagated on an extra-chromosomal plasmid. Selection of clones harboring the mismatch repair gene expression vector can be accomplished by addition of any of several different antibiotics, including but not limited
10 to ampicillin, kanamycin, chloramphenicol, zeocin, and tetracycline. The microbe can be any species for which suitable techniques are available to produce transgenic microorganisms, such as but not limited to genera including *Bacillus*, *Pseudomonas*, *Staphylococcus*, *Escherichia* and others.

Any method for making transgenic bacteria known in the art can be
15 used. According to one process of producing a transgenic microorganism, the polynucleotide is transfected into the microbe by one of the methods well known to those in the art. Next, the microbial culture is grown under conditions that select for cells in which the polynucleotide encoding the mismatch repair gene is either incorporated into the host genome as a stable
20 entity or propagated on a self-replicating extra-chromosomal plasmid, and the protein encoded by the polynucleotide fragment transcribed and subsequently translated into a functional protein within the cell. Once transgenic microbe is engineered to harbor the expression construct, it is then propagated to generate and sustain a culture of transgenic microbes
25 indefinitely.

Once a stable, transgenic microorganism has been engineered to express a functional mismatch repair (MMR) protein, the microbe can be exploited to create novel mutations in one or more target gene(s) of interest harbored within the same microorganism. A gene of interest can be any
30 gene naturally possessed by the bacterium or one introduced into the

bacterial host by standard recombinant DNA techniques. The target gene(s) may be known prior to the selection or unknown. One advantage of employing such transgenic microbes to induce mutations in resident or extra-chromosomal genes within the microbe is that it is unnecessary to expose the microorganism to mutagenic insult, whether it be chemical or radiation in nature, to produce a series of random gene alterations in the target gene(s). This is due to the highly efficient nature and the spectrum of naturally occurring mutations that result as a consequence of the altered mismatch repair process. However, it is possible to increase the spectrum and frequency of mutations by the concomitant use of either chemicals and/or radiation together with MMR defective cells. These include DNA mutagens, DNA alkylating agents, DNA intercalating agents, DNA oxidizing agents, ionizing radiation, and ultraviolet radiation. The net effect of the combination treatment is the increase in altered gene pool in the genetically altered microbe that result in an increased alteration of an allele(s) that are useful for producing new output traits. Another benefit of using MMR-defective microbes that are taught in this application is that one can perform a genetic screen for the direct selection of variant sub-clones that exhibit new output traits with commercially important applications. This allows one to bypass tedious and time consuming gene identification, isolation and characterization.

Mutations can be detected by analyzing the recombinant microbe for alterations in the genotype and/or phenotype post-activation of the decreased mismatch repair activity of the transgenic microorganism. Novel genes that produce altered phenotypes in MMR-defective microbial cells can be discerned by any variety of molecular techniques well known to those in the art. For example, the microbial genome can be isolated and a library of restriction fragments cloned into a plasmid vector. The library can be introduced into a "normal" cell and the cells exhibiting the novel phenotype screened. A plasmid is isolated from those normal cells that

exhibit the novel phenotype and the gene(s) characterized by DNA sequence analysis. Alternatively, differential messenger RNA screen can be employed utilizing driver and tester RNA (derived from wild type and novel mutant respectively) followed by cloning the differential transcripts and characterizing them by standard molecular biology methods well known to those skilled in the art. Furthermore, if the mutant sought is on encoded by an extrachromosomal plasmid, then following co-expression of the dominant negative MMR gene and the gene of interest to be altered and phenotypic selection, the plasmid is isolated from mutant clones and analyzed by DNA sequence analysis by methods well known to those in the art. Phenotypic screening for output traits in MMR-defective mutants can be by biochemical activity and/or a physical phenotype of the altered gene product. A mutant phenotype can also be detected by identifying alterations in electrophoretic mobility, DNA binding in the case of transcription factors, spectroscopic properties such as IR, CD, X-ray crystallography or high field NMR analysis, or other physical or structural characteristics of a protein encoded by a mutant gene. It is also possible to screen for altered novel function of a protein *in situ*, in isolated form, or in model systems. One can screen for alteration of any property of the microorganism associated with the function of the gene of interest, whether the gene is known prior to the selection or unknown. The aforementioned screening and selection discussion is meant to illustrate the potential means of obtaining novel mutants with commercially valuable output traits.

Plasmid expression vectors that harbor the mismatch repair (MMR) gene inserts can be used in combination with a number of commercially available regulatory sequences to control both the temporal and quantitative biochemical expression level of the dominant negative MMR protein. The regulatory sequences can be comprised of a promoter, enhancer or promoter/enhancer combination and can be inserted either upstream or downstream of the MMR gene to control the expression level. The

regulatory promoter sequence can be any of those well known to those in the art, including but not limited to the lacI, tetracycline, tryptophan-inducible, phosphate inducible, T7-polymerase-inducible (30), and steroid inducible constructs as well as sequences which can result in the excision of the dominant negative mismatch repair gene such as those of the Cre-Lox system. These types of regulatory systems are familiar to those skilled in the art.

Once a microorganism with a novel, desired output trait of interest is created, the activity of the aberrant MMR activity can be attenuated or eliminated by any of a variety of methods, including removal of the inducer from the culture medium that is responsible for promoter activation, gene disruption of the aberrant MMR gene constructs, electroporation and/or chemical curing of the expression plasmids (Brosius, Biotechnology 10:205-225, 1988; Wang et al., J. of Fujian Agricultural University 28:43-46, 1999; Fu et. al., Chem Abstracts 34:415-418, 1988). The resulting microbe is now useful as a stable strain that can be applied to various commercial applications, depending upon the selection process placed upon it.

In cases where genetically deficient mismatch repair bacteria [strains such as but not limited to: M1 (mutS) and in EC2416 (mutS delta umuDC), and mutL or mutY strains] are used to derive new output traits, transgenic constructs can be used that express wild-type mismatch repair genes sufficient to complement the genetic defect and therefore restore mismatch repair activity of the host after trait selection [Grzesiuk, E. et.al. (Mutagenesis 13:127-132, 1998); Bridges, B.A., et.al. (EMBO J. 16:3349-3356, 1997); LeClerc, J.E., Science 15:1208-1211, 1996); Jaworski, A. et.al. (Proc. Natl. Acad. Sci USA 92:11019-11023, 1995)]. The resulting microbe is genetically stable and can be applied to various commercial practices.

The use of over expressing foreign mismatch repair genes from human and yeast such as PMS1, MSH2, MLH1, MLH3, etc. have been previously demonstrated to produce a dominant negative mutator phenotype in bacterial hosts (35, 36, 37). In addition, the use of bacterial strains expressing prokaryotic dominant negative MMR genes as well as hosts that have genomic defects in endogenous MMR proteins have also been previously shown to result in a dominant negative mutator phenotype (29,32). However, the findings disclosed here teach the use of MMR genes, including the human PMSR2 and PMSR3 gene (ref 19), the related PMS134 truncated MMR gene (ref 32), the plant mismatch repair genes **(SARAH include Plant patent application)** and those genes that are homologous to the 134 N-terminal amino acids of the PMS2 gene which include the MutL family of MMR proteins and including the PMSR and PMS2L homologs described by Hori et.al. (accession number NM_005394 and NM_005395) and Nicolaides (reference 19) to create hypermutable microbes. In addition, this application teaches the use of DNA mutagens in combination with MMR defective microbial hosts to enhance the hypermutable production of genetic alterations. This accentuates MMR activity for generation of microorganisms with commercially relevant output traits such as but not limited to recombinant protein production strains, biotransformation, and bioremediation.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples that will be provided herein for purposes of illustration only, and are not intended to limit the scope of the invention

EXAMPLES

Example 1: Generation of inducible MMR dominant negative allele vectors

Bacterial expression constructs were prepared to determine if the human PMS2 related gene (hPMSR3) (19) and the human PMS134 gene (32) are capable of inactivating the bacterial MMR activity and thereby increase the overall frequency of genomic hypermutation, a consequence of which is the generation of variant sib cells with novel output traits following host selection. Moreover, the use of regulatable expression vectors will allow for suppression of dominant negative MMR alleles and restoration of the MMR pathway and genetic stability in hosts cells (43). For these studies, a plasmid encoding the hPMS134 cDNA was altered by polymerase chain reaction (PCR). The 5' oligonucleotide has the following structure: 5'-ACG CAT ATG GAG CGA GCT GAG AGC TCG AGT-3' that includes the NdeI restriction site CAT ATG. The 3'-oligonucleotide has the following structure: 5'-GAA TTC TTA TCA CGT AGA ATC GAG ACC GAG GAG AGG GTT AGG GAT AGG CTT ACC AGT TCC AAC CTT CGC CGA TGC-3' that includes an EcoRI site GAA TTC and the 14 amino acid epitope for the V5 antibody. The oligonucleotides were used for PCR under standard conditions that included 25 cycles of PCR (95°C for 1 minute, 55°C for 1 minute, 72°C for 1.5 minutes for 25 cycles followed by 3 minutes at 72°C). The PCR fragment was purified by gel electrophoresis and cloned into pTA2.1 (InVitrogen) by standard cloning methods (Sambrook et al., Molecular Cloning: A Laboratory Manual, Third Edition, 2001), creating the plasmid pTA2.1-hPMS134. pTA2.1-hPMS134 was digested with the restriction enzyme EcoRI to release the insert (there are two EcoRI restriction sites in the multiple cloning site of pTA2.1 that flank the insert) and the fragment filled in with Klenow fragment and dNTPs. Next, the fragment was gel purified, then digested with NdeI and inserted in pT7-Ea that had been digested with NdeI and BamHI (filled

with Klenow) and phosphatase treated. The new plasmid was designated pT7-Ea-hPMS134. The following strategy, similar to that described above to clone human PMS134, was used to construct an expression vector for the human related gene PMSR3. First, the hPMSR3 fragment was amplified
5 by PCR to introduce two restriction sites, an NdeI restriction site at the 5'-end and an EcoRI site at the 3'-end of the fragment. The 5'-oligonucleotide that was used for PCR has the following structure: 5'-ACG CAT ATG TGT CCT TGG CGG CCT AGA-3' that includes the NdeI restriction site CAT ATG. The 3'-oligonucleotide used for PCR has the
10 following structure: 5'-GAA TTC TTA TTA CGT AGA ATC GAG ACC GAG GAG AGG GTT AGG GAT AGG CTT ACC CAT GTG TGA TGT TTC AGA GCT-3' that includes an EcoRI site GAA TTC and the V5 epitope to allow for antibody detection. The plasmid that contained human PMSR3 in pBluescript SK (19) was used as the PCR target with the
15 hPMS2-specific oligonucleotides above. Following 25 cycles of PCR (95°C for 1 minute, 55°C for 1 minute, 72°C for 1.5 minutes for 25 cycles followed by 3 minutes at 72°C). The PCR fragment was purified by gel electrophoresis and cloned into pTA2.1 (InVitrogen) by standard cloning methods (Sambrook et al., Molecular Cloning: A Laboratory Manual, Third
20 Edition, 2001), creating the plasmid pTA2.1-hR3. pTA2.1-hR3 was next digested with the restriction enzyme EcoRI to release the insert (there are two EcoRI restriction sites in the multiple cloning site of pTA2.1 that flank the insert) and the fragment filled in with Klenow fragment and dNTPs. Then, the fragment was gel purified, then digested with NdeI and inserted
25 in pT7-Ea that had been digested with NdeI and BamHI (filled with Klenow) and phosphatase treated. The new plasmid was designated pT7-Ea-hR3.

BL21 cells harbor an additional expression vector for the lysozyme protein, which has been demonstrated to bind to the T7 polymerase in situ;
30 this results in a bacterial strain that has very low levels of T7 polymerase

expression. However, upon addition of the inducer IPTG, the cells express high-levels of T7 polymerase due to the IPTG-inducible element that drives expression of the polymerase that is resident within the genome of the BL21 cells (30). The BL21 cells are chloramphenicol resistant due to the plasmid that expresses lysozyme within the cell. To introduce the pT7-hPMS134 or the pT7-hPMSR3 genes into BL21 cells, the cells were made competent by incubating the cells in ice cold 50mM CaCl₂ for 20 minutes, followed by concentrating the cells and adding super-coiled plasmid DNA as describe (Maniatis, T.et.al. Cold Spring Harbor Laboratory Press, Third Edition, 2001). Ampicillin resistant BL21 were selected on LB-agar plates [5% yeast extract, 10% bactotryptone, 5% NaCl, 1.5% bactoagar , pH 7.0 (Difco)] plates containing 25 µg/ml chloramphenicol and 100µg/ml ampicillin. The next day, bacterial colonies were picked and analyzed for vectors containing an intact pTACPMS134 or pTAC empty vector by restriction endonuclease digestion and sequence analysis.

In addition to constructing a V5-epitope tagged PMS134 construct we also constructed and tested a non-epitope tagged version. This was prepared to demonstrate that the simple fact of epitope tagging the construct did not result in alteration of the dominant-negative phenotype that PMS134 has on mismatch repair activity. For these studies, a BamHI restriction fragment containing the hPMS134 cDNA was filled-in with Klenow fragment and then sub-cloned into a Klenow-filled blunt-ended NdeI-XhoI site of the pTACLAC expression vector, which contains the isopropylthio-β-galactosidase (IPTG)-inducible bacterial TAC promoter and ampicillin resistance gene as selectable marker. The NdeI-XhoI cloning site is flanked by the TACLAC promoter that contains the LacI repressor site followed by a Shine Dalgarno ribosome-binding site at the 5' flanking region and the T1T2 ribosomal RNA terminator in the 3' flanking region. The TACLAC vector also contains the LacI gene, which is constitutively expressed by the TAC promoter.

5 DH10B bacterial cells containing the pBCSK vector (Stratagene), which constitutively expresses the β -galactosidase gene and contains the chloramphenicol resistance marker for selection, were made competent via the CaCl_2 method (Maniatis, T.et.al. Cold Spring Harbor Laboratory Press, 1982). This vector turns bacterial cells blue when grown in the presence of IPTG and X-gal that aids in the detection of bacterial colonies. Competent cells were transfected with the pTAC empty vector or the pTACPMS134 vector following the heat-shock protocol. Transfected cultures were plated onto LB-agar [5% yeast extract, 10% bactotryptone, 5% NaCl, 1.5% bactoagar , pH 7.0 (Difco)] plates containing 25 $\mu\text{g/ml}$ chloramphenicol and 100 $\mu\text{g/ml}$ ampicillin. The next day, bacterial colonies were picked and analyzed for vectors containing an intact pTACPMS134 or pTAC empty vector by restriction endonuclease digestion and sequence analysis. Ten clones of each bacteria containing correct empty or PMS134 inserts were then grown to confluence overnight in LB media (5% yeast extract, 10% bactotryptone, 5% NaCl, pH 7.0) containing 10 $\mu\text{g/ml}$ chloramphenicol and 50 $\mu\text{g/ml}$ ampicillin. The next day TAC empty or pTACPMS134 cultures were diluted 1:4 in LB medium plus 50 μM IPTG (Gold Biotechnology) and cultures were grown for 12 and 24 hours at 37°C. After incubation, 50 μl aliquots were taken from each culture and added to 150 μl s of 2X SDS buffer and cultures were analyzed for PMS134 protein expression by western blot.

Western blots were carried out as follows. 50 μl s of each PMS134 or empty vector culture was directly lysed in 2X lysis buffer (60 mM Tris, pH 6.8, 2% SDS, 10% glycerol, 0.1 M 2-mercaptoethanol, 0.001% bromophenol blue) and samples were boiled for 5 minutes. Lysate proteins were separated by electrophoresis on 4-20% Tris glycine gels (Novex). Gels were electroblotted onto Immobilon-P (Millipore) in 48 mM Tris base, 40 mM glycine, 0.0375% SDS, 20% methanol and blocked overnight

at 4°C in Tris-buffered saline plus 0.05% Tween-20 and 5% condensed milk. Filters were probed with a rabbit polyclonal antibody generated against the N-terminus of the human PMS2 polypeptide (Santa Cruz), which is able to recognize the PMS134 polypeptide (31), followed by a secondary goat anti-rabbit horseradish peroxidase-conjugated antibody. After incubation with the secondary antibody, blots are developed using chemiluminescence (Pierce) and exposed to film to measure PMS134 expression.

As shown in Figure 1, a robust expression of PMS134 could be detected in bacterial cells containing pTACPMS134 (lane 2) in contrast to cells expressing empty vector (lane 1), which had no signal.

For induction of PMS134 and PMSR3 in BL21 cells, the pT7-Ea-hPMS134 or the pT7-Ea-hPMSR3 cells were induced with 50µM IPTG for 12 and 24 hours. Cell lysates were prepared and analyzed by western blot listed above using either the N-terminal PMS2 antibody to detect the PMS134 containing cells or the antiV5-horseradish peroxidase conjugated monoclonal antibody (InVitrogen) to detect the PMS134V5 and PMSR3V5 polypeptides. Figure 2 shows the expression of PMS134V5 and PMSR3V5 before (-) lanes and after IPTG (+) lanes induction.

Example 2: Generation of hypermutable bacteria with inducible dominant negative alleles of mismatch repair genes

Bacterial clones expressing the *PMS134* or the empty vector were grown in liquid culture for 24 hr at 37°C in the presence of 10 µg/ml chloramphenicol and 50 µg/ml ampicillin plus 50µM IPTG. The next day, cultures were diluted 1:10 in medium containing 50µM IPTG plus ampicillin/chloramphenicol (AC) or ampicillin/chloramphenicol plus 25 µg/ml kanamycin (ACK) and cultures were grown for 18 hr at 37°C. The following day, a 0.1 µl aliquot (2 µl diluted in 1000 µl of LB medium and

used 50 µl for plating) of cells grown in AC medium were plated on LB-
agar plates containing 40 µg/ml of 5-bromo-4-chloro-3-indolyl-B-D-
galactoside (X-gal) plus 100µg/ml ampicillin (AMP), while a 1 µl aliquot
(1 µl diluted in 100 µl of LB medium and used 100 µl for plating) of cells
5 grown in ACK medium were plated on LB-agar plates containing X-gal
and 50µg/ml kanamycin (KAN). Plates were incubated for 18 hours at
37°C. The results from these studies show that cells expressing the
PMS134 were able to increase hypermutation in the genome of the DH10B
bacterial strain which resulted in the production of siblings that exhibit new
10 biological traits such as KAN resistance (Figure 3).

Kanamycin-resistant assays using BL21 cells expressing the
V5-tagged or untagged PMS134 or PMSR3 polypeptides were
carried out as described above. BL21 bacterial cells that harbor the
empty vector, pT7-PMS134 or pT7-PMSR3 were grown overnight in
15 LB supplemented with 100ug/ml ampicillin. The overnight cultures
were diluted 1:100 into fresh ampicillin containing medium and
grown for 2.5 hours at 37°C with continuous shaking. When the cells
reached an optical density (OD) of 0.6, measured at 600nm, IPTG
was added to each culture to a final concentration of 0.5mM. Cells
20 were incubated for 24, and 48 hours; at those time points cells were
removed for SDS-PAGE analysis and plating (see above).
BL21/pT7 (empty vector), BL21/pT7-PMS134, and BL21/pT7-R3
cells were plated onto LB plates, LB plates that contained 100ug/ml
ampicillin, and plates that contain 50ug/ml Kanamycin. The
25 equivalent of 1×10^7 cells/plate were spread onto the plates. BL21
cells that harbor the empty vector are capable of growth on LB plates
as well as LB plates that contain 100ug/ml ampicillin; that is as
expected since the pT7 expression vector renders the cells ampicillin
resistant. The vector only control is not capable of growth on
30 Kanamycin. After 24hr IPTG-induction PMS134 or PMSR3 cells

had a significant number of KAN resistant cells while none were observed in BL21 parental cells grown under similar conditions (Figure 4). Moreover, BL21 cells containing the PMS134 or PMSR3 genes under non-IPTG-induced conditions failed to produce any KAN resistant clones demonstrating the need for expression of the PMS polypeptides for hypermutability. A summary outlining the data and number of Kanamycin resistant bacterial clones is provided in TABLE 1.

10 **TABLE 1. Generation of Kanamycin resistant clones via MMR deficiency**

| STRAIN | # CELLS SEEDED | AMP ^R colonies | KAN ^R colonies | FREQUENCY |
|---------------------|----------------|---------------------------|---------------------------|----------------------|
| DH10B VEC | 50,000 | 62,000 | 0 | 0 |
| DH10B PMS134 | 50,000 | 43,146 | 23 | 5.3×10^{-4} |
| BL21 VEC | 500,000 | 520,80 0 | 0 | 0 |
| BL21 T7-Ea-PMS134V5 | 500,000 | 450,00 0 | 2,245 | 4.9×10^{-3} |
| BL21 T7-Ea-PMSR3V5 | 500,000 | 500,00 0 | 1,535 | 3.1×10^{-3} |

These data demonstrate and enable the proof-of-concept that the use of the dominant negative MMR genes is a viable approach to creating hypermutable bacteria that can lead to the generation of phenotypically diverse offspring when put under selective conditions.

Using the same protocol as listed above and the same cloning strategy, a truncated PMS2 homolog from the *Arabidopsis thaliana* plant (**reference the PLANT patent**), which was cloned by degenerate PCR from an *Arabidopsis thaliana* cDNA library

(Stratagene), was found to give a similar enhancement of genetic hypermutability in DH5alpha bacteria Figure 5. For detection purposes, we fused a FLAG epitope to the C-terminus of the PMS134 polypeptide using PCR and an antisense primer directed to the 134 codon region of the *Arabidopsis* PMS2 homolog followed by a FLAG epitope and 2 termination codons. The resultant fusion was termed ATPMS134-flag. The AT PMS134-flag gene was then cloned into the IPTG-inducible TACLAC expression vector and transfected into DH5alpha cells. Western blot of bacteria transfected with an IPTG-inducible expression vector carrying a truncated version (codons 1-134) of the *Arabidopsis thaliana* PMS2 homolog using the anti-FLAG antibody demonstrated the inducibility and steady-state protein levels of the chimeric gene. Figure 5A shows the western blot containing protein from an untransfected cell (lane 1) and a bacterial clone expressing the *Arabidopsis* PMS2-134 truncated protein (lane 2). Following the mutagenesis protocol described above, bacterial cells expressing the ATPMS134 protein were found to have an increase in the number of KAN resistant cells (12 clones) in contrast to cells expressing the empty vector that yielded no KAN resistant clone.

Bacterial cells such as the pT7-PMS134 and pT7-R3 harboring BL21 cells; the TACLACPMS134 DH10B; the TACLACMLH1 DH10B cells; or the TACLAC-ATPMS134flag DH5alpha cells are capable of growth on LB, LB/ampicillin and LB/KAN plates because the cells have acquired mutations within their genome that render the cell drug resistant. Cells that express dominant negative MMR genes have altered the mismatch control pathway of the microbe, presumably altering a gene or a set of genes that control resistance to kanamycin. A new output trait, Kanamycin-resistance, is generated by expression of the dominant negative MMR

gene in these cells. These data demonstrate the ability of dominant negative MMR genes to produce hypermutability across a wide array of bacterial strains to produce new output traits such as Kanamycin resistance.

5

EXAMPLE 3: Dominant negative MMR genes can produce new genetic variants and commercially viable output traits in prokaryotic organisms.

10 The data presented in EXAMPLE 2 show the ability to generate genetic alterations and new phenotypes in bacterial strains expressing dominant negative MMR genes. In this EXAMPLE we teach the utility of this method to create prokaryotic strains with commercially relevant output traits.

15

GENERATION OF HEAT-RESISTANT PRODUCER STRAINS.

One example of commercial utility is the generation of heat-resistant recombinant protein producer strains. In the scalable
20 process of recombinant manufacturing, large-scale fermentation of prokaryotes results in the generation of heat, which leads to suboptimal growth conditions for the producer strain and thus resulting in lower recombinant protein yields. In order to circumvent this problem, we employed the use of DH10B bacteria containing the
25 inducible TACLACPMS134 gene. Briefly, cells were grown in 5 ml LB shake flasks containing ampicillin and IPTG-induced for 0, 24 and 48hrs at 37C. Cultures were harvested and then incubated at 100C for 0, 1 or 10 minutes (times at which 100% of the wild-type strain perishes) and 100 µl aliquots (equivalent to 250,000 cells) were
30 plated onto LB agar plates containing ampicillin to identify heat

resistant clones. Table 2 shows a typical experiment whereby cells containing the TACLACPMS134 gene generated a significant number of heat-resistant clones after 48 hours of PMS134 induction and hypermutation via MMR blockade. No or a few clones were observed in the uninduced or 24hr induced conditions respectively suggesting the needs for multiple rounds of genetic mutation to produce genes that are capable of allowing bacteria to survive under harsh conditions. Similar results were observed with other dominant negative mutants such as the PMSR2, PMSR3, and the human MLH1 proteins (not shown).

TABLE 2. Generation of heat-resistant clones via MMR deficiency

| Treatment | Heated 0 min | Heated 1 min | Heated 10 min |
|----------------------------|-----------------------|-----------------|------------------|
| TACLACVEC 0 hr IPTG | 250,000 +/- 7,500 | 0 | 0 |
| TACLACPMS134 0 hr IPTG | 265,000 +/- 2,000 | 0 | 0 |
| TACLACVEC 24 hr IPTG | 274,000 +/- 12,000 | 1 +/- 0 | 0 |
| TACLACPMS134 24 hr IPTG | 240,000 +/- 9,400 | 5 +/- 2 | 0 |
| TACLACVEC 48 hr IPTG | 256,000 +/- 12,000 | 0 | 0 |
| TACLACPMS134 48 hr IPTG | 252,000 +/- 14,000 | 65 +/- 8 | 3 +/- 1 |

15 GENERATION OF HIGH RECOMBINANT PROTEIN PRODUCER STRAINS.

Next, we tested the ability of bacteria expressing dominant negative MMR genes to produce subclones with enhanced recombinant protein production. In these experiments again we employed the DH10B cells containing the TACLACPMS134 inducible vector plus the pTLACZ vector, which constitutively

expresses the β -galactosidase gene. Analysis of individual clones containing the TACLACPMS134 and pTLACZ vector typically produces 10-20 μ g/ml of LACZ protein via shake flask fermentation after IPTG induction for 24 hours. To test the hypothesis that high recombinant producer strains can be generated by decreased MMR in bacterial strains, we induced the TACLACPMS134-pTLACZ cells for 48 hours with IPTG as described above. We then diluted the culture 1:50 in LB medium, grew the strain for 24 hours, and plated 10 μ ls of culture (diluted in 300 μ ls of LB) onto LB amp-XGAL plates to identify candidate clones that produce robust levels of recombinant LACZ protein. As a control, uninduced cells were treated similarly and plated onto LB amp-XGAL plates. Analysis of the plates revealed a number of bacterial colonies exhibiting a number of clones with an intense BLUE staining in the TACLACPMS134/pTLACZ cells induced with IPTG but none were observed in uninduced clones (Figure 6). To confirm that these clones produced an enhanced level of LACZ, we expanded 2 clones with an average BLUE stain (BGAL-C1 and BGAL-C2) and 10 clones with a robust BLUE staining (BGAL-MOR1 to BGAL-MOR10). We grew all clones in LB AMP for 24 hours without IPTG and replated the clones. Six out of ten BGAL-MOR clones resulted in a more robust β -gal stain *in situ* as compared to control "average" cells (BGALC1 and C2). We next performed a more quantitative assay using a β -gal ELISA assay. Briefly, 2 mls of cell centrifuged at 10,000gs for 10 minutes and resuspended in 0.5 mls of 0.25M Tris, pH 7.5 plus 0.0001% Tween-20. Cells were freeze-thawed 4X's and vortexed for 4 minutes at room temperature. Lysates were cleared of debris by centrifugation and supernatants were collected. Protein extracts were quantified for total protein using the Bradford assay (BioRad) as described by the manufacturer. Plate ELISAs were

carried out by coating 96 well maxisorb (NUNC) plates with 0.1 mls of a 1 µg/ml (diluted in PBS pH7.0) bacterial extract solution and a dose range of recombinant β-GAL (Sigma) from 0.001 to 10 mg/ml. All samples were plated in triplicates. Plates were coated for 2 hours, washed 2 times with PBS and blocked with 0.2mls of PBS plus 5% powdered milk for 30 minutes. Next, plates were washed once with PBS and incubated with an anti-β-galactosidase monoclonal antibody that recognizes both native and denatured forms (Sigma) for 2 hours. Plates were then washed 3 times with PBS and incubated with 0.1 mls of an anti-mouse horseradish peroxidase conjugated antibody for 1 hour at room temperature. Plates were washed 3 times with PBS and incubated with TMB ELISA substrate (BioRad) for 15 to 30 minutes. Reactions were stopped with 0.1N H₂SO₄ and read on a BioRAD plate reader at 415nm. The control clones produced roughly 9 and 13 µgs /ml of β-gal while BGAL-MOR clones 2, 3 and 9 produced 106, 82 and 143 µgs /ml of β-gal. To determine if reason that these clones produced more β-gal was due to mutations in the plasmid promoter elements, we isolated the pTLACZ plasmid and retransfected it into DH10B cells as described above. *In situ* analysis found the resultant clones to produce similar amount of β-gal as that of the control. These data suggest that the BGAL-MOR 2, 3, and 9 hosts had alterations, which results in elevated expression and/or stability of recombinant proteins.

To determine if the enhanced *in situ* β-gal expression that was observed in BGAL-MOR clones 1, 5, and 6, which did not appear to have enhanced β-gal protein levels (had less than 15 µg/ml as determined by ELISA) was authentic, we performed a more quantitative assay on these lines plus the BGAL-MOR 9, the BGALC1 and C2 lines as control. Cells containing an empty vector

(without a LACZ gene) were used as negative control. To measure β -gal activity, we employed a colorimetric β -gal substrate assay using CPRG (Roche) as described (31). Briefly, 5 μ gs of protein extract isolated for ELISA analysis (described above) were analyzed using a plate assay. Protein was added to buffer containing 45 mM 2-mercaptoethanol, 1mM MgCl₂, 0.1 M NaPO₄ and 0.6 mg/ml Chlorophenol red- β -D-galactopyranoside (CPRG, Roche). Reactions were incubated for 1 hour, terminated by the addition of 0.5 M Na₂CO₃, and analyzed by spectrophotometry at 576 nm in a BioRad plate reader. Analysis of these extracts confirmed our *in situ* data that these cells did have increased β -gal activity (TABLE 3).

TABLE 3. Generation of bacterial clones with increased β -gal enzymatic activity via MMR deficiency.

| Clone | β -gal protein (μ g/ml) | β -gal activity (O.D. 576) |
|--------------|------------------------------------|----------------------------------|
| BGAL-C1 | 9 | 0.413 +/- .092 |
| BGAL-C2 | 13 | 0.393 +/- .105 |
| BGAL-MOR1 | 14 | 0.899 +/- .134 |
| BGAL-MOR5 | 13 | 0.952 +/- .133 |
| BGAL-MOR6 | 15 | 0.923 +/- .100 |
| BGAL-MOR9 | 143 | 0.987 +/- .106 |
| Empty vector | - | 0.132 +/- .036 |

Because there was no observable increase in the amount of β -gal protein one likely hypothesis is that the β -gal gene structure was mutated during the hypermutability growth stage and now produces a

more active enzyme. Sequence analysis confirms that this may be the reason for enhanced activity in a subset of clones.

Together, these data demonstrate the ability to produce genetically altered prokaryotic host strains using dominant negative
5 MMR genes to generate commercially valuable output traits such as high recombinant protein producer lines and structurally altered enzymes with enhanced activities.

**EXAMPLE 4: Mutations in the host genome generated by
10 defective MMR are genetically stable**

As described in EXAMPLE 2 and 3, manipulation of the MMR pathway in microbes results in alterations within the host genome and the ability to select for a novel output traits. It is
15 important that the mutations introduced as a result of defective MMR is genetically stable and passed on to daughter cells once a desired output pathway is established. To determine the genetic stability of mutations introduced into the microbial genome the following experiment was performed. Five independent colonies from pT7-
20 PMS134 and pT7-PMSR3 that are kanamycin resistance were grown overnight from an isolated colony in 5 ml of LB. Next, 1 μ L of the overnight culture from these cultures were inoculated into another 5 mL of LB and grown overnight to saturation. Under these growth conditions the microbial cells have divided over 20 generations.
25 Therefore, if the new output trait generated by alteration of MMR is unstable, the cells should “revert” back from kanamycin resistance to kanamycin sensitivity. Cells were plated onto LB plates and incubated overnight at 37⁰C. Next, the colonies (about 1,000/plate) were replica plated to LB, LB^{amp100}, and LB^{kan50} plates and incubated
30 at 37⁰C overnight. Analysis of clones from these studies reveal that a

strict correlation occurs with loss of dominant negative MMR expression and phenotype stability. No loss of KAN resistant clones generated in Example 3 were observed when cells were grown in the absence of IPTG (not expressing PMS134), while 5 revertants out of 1200 were observed in clones the were continually grown in IPTG (express PMS134). Extended culturing of cells and replica plating found no reversions of KAN resistance in cultures grown in the absence of IPTG, which produce no PMS134 as determined by western blot (data not shown).

These data demonstrate the utility of employing inducible expression systems and dominant negative MMR genes in prokaryotes to generate genetically altered strains for commercial applications such as but not limited to enhanced recombinant manufacturing and biotransformation that can then in turn be restored to a genetically stable host with a “fixed” new genotype that is suitable for commercial processes.

EXAMPLE 5: Enhanced Generation of MMR-Defective Bacteria and Chemical Mutagens for the Generation of New Output

Traits

It has been previously documented that MMR deficiency yields to increased mutation frequency and increased resistance to toxic effects of chemical mutagens (CM) and their respective analogues such as but not limited to those as: ethidium bromide, EMS, MNNG, MNU, Tamoxifen, 8-Hydroxyguanine, as well as others listed but not limited to in publications by: Khromov-Borisov, N.N., et.al. (Mutat. Res. 430:55-74, 1999); Ohe, T., et.al. (Mutat. Res. 429:189-199, 1999); Hour, T.C. et.al. (Food Chem. Toxicol. 37:569-579, 1999); Hrelia, P., et.al. (Chem. Biol. Interact. 118:99-111, 1999); Garganta, F., et.al. (Environ. Mol. Mutagen. 33:75-85, 1999);

Ukawa-Ishikawa S., et.al. (Mutat. Res. 412:99-107, 1998);
www.ehs.utah.edu/ohh/mutagens, etc. To demonstrate the ability of CMs
to increase the mutation frequency in MMR defective bacterial cells, we
exposed T7-PMS134 BL21 cells to CMs.

5 T7-PMS134 cells and empty vector control cells were grown with
IPTG for 48 hours and then diluted 1: 50 in LB plus IPTG and increasing
amounts of ethyl methane sulfonate (EMS) from 0, 1, 10, 50, 100, and 200
 μ M. 10 μ L aliquots of culture (diluted in 300 μ l LB) were plated out on
LB agar plus ampicillin plates and grown overnight at 37C. The next day
10 plates were analyzed for cell viability as determined by colony formation.
Analysis found that while no significant difference in colony number was
observed between the pT7-PMS134 and control at the 0, 1 , or 10 μ M
concentrations (all had > 1000 colonies), the number of control cells were
reduced to 30 and 0 at the 50 and 100 μ M concentrations, respectively. No
15 difference was observed in the pT7-PMS134 cells treated with 0, 1, 10 or
50 μ M, while a 3 fold reduction was observed in cultures treated with 100
 μ M EMS. The 200 μ M treatment was toxic for both lines. These data
demonstrate the ability of MMR deficiency to protect prokaryotes against
the toxic effects of DNA alkylating agents and provides a means to generate
20 a wider range of mutations that can lead to an increased number of genetic
variations and an increase in the number of new biochemical activities
within host proteins to produce new output traits for commercial
applications.

To confirm that MMR deficient bacterial cells treated with CM can
25 result in an increased mutation rate and produce a greater number of
variants, we cultured pT7-PMS134 cells and empty vector controls in the
presence of IPTG for 48 hours, followed by dilution and regrowth in 25 μ M
EMS for 24 hours as described above. Cells were plated out on 100 mM
petri dishes containing ampicillin or KAN and scored for KAN resistance.

30 Analysis revealed that an 11-fold increase in the generation of KAN

resistant cells were found in pT7-Ea-PMS134V5 cells in contrast to control cells.

These data demonstrate the use of employing a regulated dominant negative MMR system plus chemical mutagens to produce enhanced numbers of genetically altered prokaryotic strains that can be selected for new output traits. This methods is now useful generating such organisms for commercial applications such as but not limited to recombinant manufacturing, biotransformation, and altered biochemicals (biotransformation) with enhanced activities for manufacturing purposes and gene discovery for pharmaceutical compound development.

EXAMPLE 6: Alternative Methods to Inhibition of Bacterial MMR Activity

The inhibition of MMR activity in a host organism can be achieved by introducing a dominant negative allele as shown in EXAMPLES 2 and 3. This application also teaches us the use of using regulated systems to control MMR in prokaryotes to generate genetic diversity and output traits for commercial applications. Other ways to regulate the suppression of MMR activity of a host is by using genetic recombination to knock out alleles of a MMR gene that can be spliced out such after selection using a system such as the CRE-Lox system; 2) blocking MMR protein dimerization with other subunits (which is required for activity) by the introduction of polypeptides or antibodies into the host via transfection methods routinely used by those skilled in the art; or 3) decreasing the expression of a MMR gene using anti-sense oligonucleotides.

MMR gene knockouts. We intend to generate disrupted targeting vectors of a particular MMR gene and introduce it into the genome of bacteria using methods standard in the art. Bacteria exhibiting hypermutability will be useful to produce genetically diverse offspring for commercial applications.

Bacteria will be confirmed to have lost the expression of the MMR gene using standard northern and biochemical techniques (as described in reference 32). MMR gene loci can be knocked out, strains selected for new output traits and MMR restored by introducing a wildtype MMR gene to complement the KO locus. Other strategies include using KO vectors that can target a MMR gene locus, select for host output traits and then have the KO vector “spliced” from the genome after strain generation. This process could be performed using systems such as but not limited to CRE-Lox.

- 10 Blocking peptides. MMR subunits (MutS and MutL proteins) interact to form active MMR complexes. Peptides are able to specifically inhibit the binding of two proteins by competitive inhibition. The use of peptides or antibodies to conserved domains of a particular MMR gene can be introduced into prokaryotic cells using lipid transfer methods that are standard in the art. Bacteria will be confirmed to have lost the expression of the MMR gene using standard northern and biochemical techniques (as described in reference 32). Bacteria exhibiting hypermutability will be useful to produce genetically diverse sibs for commercial applications.

20 Discussion

- The results described above will lead to several conclusions. The expression of dominant negative MMR proteins results in an increase in hypermutability in bacteria. This activity is due to the inhibition of MMR biochemical activity in these hosts. This method provides a claim for use of dominant negative MMR genes and their encoded products for the creation of hypermutable bacteria to produce new output traits for commercial applications.

EXAMPLES OF MMR GENES AND ENCODED POLYPEPTIDES

Yeast MLH1 cDNA (accession number U07187)

```

5      1 aaataggaat gtgatacctt ctattgcatg caaagatagt gtaggaggcg ctgctattgc
      61 caaagacttt tgagaccgct tgctgtttca ttatagtgtg ggagttctcg aagacgagaa
     121 attagcagtt ttcggtgttt agtaatcgcg ctagcatgct aggacaattt aactgcaaaa
     181 ttttgatacg atagtgatag taaatggaag gtaaaaataa catagacctt tcaataagca
     241 atgtctctca gaataaaagc acttgatgca tcagtgggta acaaaattgc tgcaggtgag
    10  301 atcataatat cccccgtaaa tgctctcaaa gaaatgatgg agaattccat cgatgcgaat
     361 gctacaatga ttgatattct agtcaaggaa ggaggaatta aggtacttca aataacagat
     421 aacggatctg gaattaataa agcagacctg ccaatcttat gtgagcgatt cacgacgtcc
     481 aaattacaaa aattcgaaga tttaggtcag attcaaacgt atggattccg aggagaagct
     541 tttagccagta tctcacatgt ggcaagatgc acagtaacga caaaagttaa agaagacaga
    15  601 tgtgcatgga gagtttcata tgcagaaggt aagatgtttg aaagcccaa acctgtgtgt
     661 ggaaaagacg gtaccacgat cctagttgaa gacctttttt tcaatattcc ttctagatta
     721 agggccttga ggtcccataa tgatgaatac tctaaaatat tagatgttgt cgggcgatac
     781 gccattcatt ccaaggacat tggcttttct tgtaaaaagt tcggagactc taattattct
     841 ttatcagtta aaccttcata tacagtccag gataggatta ggactgtgtt caataaatct
    20  901 gtggccttga atttaattac ttttcatatc agcaaaagtag aagatttaaa cctggaaagc
     961 gttgatggaa aggtgtgtga tttgaatttc atatcaaaa agtccatttc attaattttt
    1021 ttcattaata atagactagt gacatgtgat cttctaagaa gagctttgaa cagcgtttac
    1081 tccaattatc tgccaagggt cttcagacct tttatttatt tgggaattgt tatagatccg
    1141 gcggtctgtg atgttaacgt tcacccgaca aagagagagg ttctgttccg gagccaagat
    25  1201 gagatcatag agaaaatcgc caatcaattg cagcgcgaat tatctgccat tgatacttca
     1261 cgtactttta aggccttctc aatttcaaca aacaagccag agtcattgat accatttaat
     1321 gacaccatag aaagtgatag gaataggaag agtctccgac aagcccaagt ggtagagaat
     1381 tcatatacga cagcaaatag tcaactaagg aaagcgaaaa gacaagagaa taaactagtc
     1441 agaatagatg cttcacaagc taaaattacg tcatttttat cctcaagtca acagttcaac
    30  1501 tttgaaggat cgtctacaaa ggcacaactg agtgaaccca aggtaaacaa tgtaagccac
     1561 tcccaagagg cagaaaagct gacactaaat gaaagcgaac aaccgcgtga tgccaatata
     1621 atcaatgata atgacttgaa ggatcaacct aagaagaaac aaaagttggg ggattataaa
     1681 gttccaagca ttgccgatga cgaagaagat gcactcccg a ttcaaaaga cgggtatatt
     1741 agagtaccta agggagcagat taatgttaat cttacagagta tcaagaaatt gcgtgaaaaa
    35  1801 gtagatgatt cgatacatcg agaactaaca gacatttttg caaatttgaa ttacgttggg
     1861 gttgtagatg aggaaagaag attagccgct attcagcatg acttaaaagct ttttttaata
     1921 gattacggat ctgtgtgcta tgagctattc tatcagattg gtttgacaga cttcgcaaac
     1981 ttttgtaaga taaacctaca gagtacaaat gtgtcagatg atatagtttt gtataatctc
     2041 ctatcagaat ttgacgagtt aaatgacgat gcttccaaag aaaaaataat tagtaaaaaa
    40  2101 tgggacatga gcagtatgct aaatgagtag tattccatag aattggtgaa tgatggctca
     2161 gataatgact taaagtctgt gaagctaaaa tctctaccac tacttttaaa aggcctacatt
     2221 ccatctctgg tcaagttacc attttttata tatcgctctg gtaaagaagt tgattgggag
     2281 gatgaacaag agtgtctaga tggattttta agagagattg cattactcta tatacctgat
     2341 atgggtccga aagtcgatac actcgatgca tcgttgtcag aagacgaaaa agcccagttt
    45  2401 ataaaataga aggaacacat atcctcatta ctagaacacg ttctcttccc ttgtatcaaa
     2461 cgaaggttcc tggcccttag acacattctc aaggatgtcg tggaaatagc caaccttcca
     2521 gatctataca aagtttttga gaggtgttaa ctttaaaacg ttttggctgt aataccaaag
     2581 tttttgttta ttctctgagt gtgattgtgt ttcatittga agtgtatgcc ctttcttta
     2641 acgattcatc cgcgagattt caaaggatat gaaatattgt tgcagttagg aaagtatgtc
    50  2701 agaaatgtat attcggattg aaactcttct aatagttctg aagtcacttg gttccgtatt
     2761 gttttcgtcc tcttctcaca gcaacgattc ttgtctaagc ttattcaacg gtaccaaaag
     2821 ccgagtcctt tttatgagag aaaacatttc atcatttttc aactcaatta tcttaataatc
     2881 attttgtagt attttgaaaa caggatggta aaacgaatca cctgaatcta gaagctgtac
     2941 cttgtcccat aaaagtttta atttactgag ctttccggtc aagtaaaacta gtttatctag
    55  3001 ttttgaaccg aatattgttg gcagatttgc agtaagtcca gttagatcta ctaaaagttg
     3061 tttgacagca gccgattcca caaaaatttg gtaaaaggag atgaaagaga cctcgcgct
     3121 aatggtttgc atcaccatcg gatgtctgtt gaaaaactca ctttttgcat ggaagttatt
     3181 aacaataaga ctaatgatta ccttagaata atgtataa

```

60 Yeast MLH1 protein (accession number U07187)

```

MSLRKALDASVVKIAAGEIIISFVNALKEMMENSIDANATMI
65 DILVKEGGIKVLQITDNGSGINKADLPILCERFTTSKLQKFEDLSQIQTYGFRGEALA
SISHVARVTVTTKVKEDRCARVSYAEGKMLES PKPVAGKDGTTILVEDLFFNIPSRL
RALRSHNDEYSKILDVVGRIYAIHSDIGFSCCKFGDSNYLSVKPSYTVQDRIRTVFN
70 KSVASNLITFHISKVEDLNLESVDGKVCNLFISKKSISLIFFINNRLVTCOLLRRAL
NSVSYNLPKGRFPFIYLGIVIDPAADVNVHPTKREVRLSQDEIIKIANQLHAEL
SAIDTSRTFKASSISTNKPESLIPNDTIESDRNRKSLRQAQVENSYTTANSQRLRKA

```

5 KRQENKLVRIDASQAKITSFLSSSQFNFEGSSTKRQLSEPKVTNVSHSQEAEKLTIN
 ESEQPRDANTINDNDLKQPKKKQKLGDKVPSIADDEKNALPISKDGYIRVPKERNV
 10 VNLTSIKKLREKVDDSIHRELTDFANLNVGVVDEERRLAIIQHDCLKFLIDYGSVC
 YELFYQIGLTDIFANFGKINLQSTNVSDDIVLYNLLSEFDELNDASKEKIIISKIWDMS
 SMLNEYYSIELVNDGLDNDLKSVKLSLPLLLKGYIPSLVKLPFFIYRLGKEVDWEDE
 15 QECLDGILREIALLYIPDMVPKVDLTLDASLSEDEKAQFINRKEHISSLLEHVLFPCK
 RRFLAPRHILKDVVEIANLPDLYKVFERC

Mouse PMS2 protein

MEQTEGVSTE CAKAIKPIDG KSVHQICSGQ VILSLSTAVK ELIENSVDAG ATTIDRLRKD 60
 YGVDLIEVSD NGCGVEEENF EGLALKHHTS KIQEFADLTQ VETFGFRGEA LSSLCALSDV 120
 TISTCHGSAS VGTRLVFDHN GKITQKTPYP RPKGTTVSQV HLFYTLPVRY KEFQRNIKKE 180
 20 YSKMVQVLQA YCIISAGVRV SCTNQLGQSK RHAVVCTSGT SGMKENIGSV FGQKQLQSLI 240
 PFFVQLPPSDA VCEEYGLSTS GRHKTFSTFR ASFHSARTAP GGVQQTGSFS SSIRGPVTQQ 300
 RSLSLSMRFY HMYNRHQYPF VVLNVSDSE CVDINVTDPK RQILLQEEKL LLAVLKTSLI 360
 GMFDSANKL NVNQQLLDV EGNLVKLHTA ELEKPVPGKQ DNSPSLKSTA DEKRVASISR 420
 LREAFSLHPT KEIKSRGPET AELTRSPSE KRGVLSSYPS DVISYRGLRG SQDKLVSPDT 480
 25 SPGDCMDREK IEKDSGLSST SAGSEEEFST PEVASSFSSD YNVSSLEDRP SQETINCGDL 540
 DCRPFGTQGS LKPEDHGYQC KALPLARLSP TNAKRFKTEE RPSNVNISQR LPGPQSTSAA 600
 EVDVAIKMNK RIVLLEFSL SAKRMKQLQ HLKAQNKHEL SYRKFRKIC PGENQAAEDE 660
 LRKEISKSMF AEMEILGQFN LGFIYTKLKE DLFLVDQHAA DEKYNFEMLQ QHTVLQAQRL 720
 ITPQTLNLTA VNEAVLIENL EIFRKNQFDF VIDEAPVTE RAKLISLPTS KNWTFGPQDI 780
 30 DELIFMLSDS PGVMCRPSRV RQMFASRACR KSMVIGTALN ASEMKKLITH MGEMDHPWNC 840
 PHGRPTMRHV ANLDVISQN
 859

Mouse PMS2 cDNA

35 gaattccggt gaaggtcctg aagaatttcc agattcctga gtatcattgg aggagacaga 60
 taacctgtcg tcaggtaacg atggtgtata tgcaacagaa atgggtgttc ctggagacgc 120
 gtcttttccc gagagcggca ccgcaactct cccgcggtga ctgtgactgg aggagtcctg 180
 catccatgga gcaaaccgaa ggcgtgagta cagaatgtgc taaggccatc aagcctattg 240
 atgggaagtc agtccatcaa atttgttctg ggcaggtgat actcagttta agcaccgctg 300
 40 tgaaggagtt gatagaaaat agtgtagatg ctggtgctac tactattgat ctaaggctta 360
 aagactatgg ggtggacctc attgaagttt cagacaatgg atgtggggta gaagaagaaa 420
 actttgaagg tctagctctg aaacatcaca catctaagat tcaagagttt gccgacctca 480
 cgcaggttga aactttcggc tttcgggggg aagctctgag ctctctgtgt gcactaagtg 540
 atgtcactat atctacctgc caccgggtctg caagcgttgg gactcgactg gtgtttgacc 600
 45 ataattggaa aatcacccag aaaactccct acccccgacc taaaggaaac acagtcagtg 660
 tgcagcactt attttataca ctaccctgct gttacaaaga gtttcagagg aacattaaaa 720
 aggagtattc caaaatggtg caggtcttac aggcgtactg tatcatctca gcaggcgtcc 780
 gtgtaagctg cactaatcag ctccgacagg ggaagcggca cgctgtggtg tgcacaagcg 840
 gcacgtctgg catgaaggaa aatatcgggt ctgtgtttgg ccagaagcag ttgcaaagcc 900
 50 tcattccttt tgttcagctg cccctagtg acgctgtgtg tgaagagtac ggcctgagca 960
 cttcaggacg ccacaaaacc tttctacgt ttcgggcttc atttcacagt gcacgcacgg 1020
 cgccgggagg agtgcaacag acaggcagtt tttcttcac aatcagaggc cctgtgaccc 1080
 agcaaaggtc tctaagcttg tcaatgaggt tttatcacat gtataaccgg catcagtacc 1140
 catttgcgt ccttaacgtt tccgttgact cagaatgtgt ggatattaat gtaactccag 1200
 55 ataaaaggca aattctacta caagaagaga agctattgct ggccgtttta aagacctcct 1260
 tgataggaat gtttgacagt gatgcaaca agcttaatgt caaccagcag ccactgctag 1320

atgttgaagg taacttagta aagctgcata ctgcagaact agaaaagcct gtgccaggaa 1380
agcaagataa ctctccttca ctgaagagca cagcagacga gaaaagggtg gcatccatct 1440
ccaggctgag agaggccttt tctcttcac cactaaaga gatcaagtct aggggtccag 1500
5 agactgctga actgacacgg agttttccaa gtgagaaaag gggcgtgtta tcctcttacc 1560
cttcagacgt catctcttac agaggcctcc gtggctcgca ggacaaattg gtgagtccca 1620
cggacagccc tgggtactgt atggacagag agaaaataga aaaagactca gggctcagca 1680
gcacctcagc tggctctgag gaagagttca gcaccccgca agtggccagt agcttttagca 1740
gtgactataa cgtgagctcc ctagaagaca gaccttctca ggaaccata aactgtgggtg 1800
10 acctggactg ccgtctctca ggtacaggac agtccttgaa gccagaagac catggatata 1860
aatgcaaaag tctacctcta gctcgtctgt caccacaaa tgccaagcgc ttcaagacag 1920
aggaaagacc ctcaaatgtc aacatttctc aaagattgcc tggctctcag agcacctcag 1980
cagctgaggt cgatgtagcc ataaaaatga ataagagaat cgtgctcctc gaggctctc 2040
tgagttctct agctaagcga atgaagcagt tacagcacct aaaggcgcag aacaaacatg 2100
aactgagtta cagaaaattt agggccaaga tttgccctgg agaaaaccaa cgagcagaag 2160
15 atgaactcag aaaagagatt agtaaatcga tgtttgcaga gatggagatc ttgggtcagt 2220
ttaacctggg atttatagta accaaactga aagaggacct ctctctgggtg gaccagcatg 2280
ctgcggatga gaagtacaac tttgagatgc tgcagcagca cagggtgctc caggcgcaga 2340
ggctcatcac accccagact ctgaacttaa ctgctgtcaa tgaagctgta ctgatagaaa 2400
atctggaaat attcagaaa aatggctttg actttgtcat tgatgaggat gctccagtca 2460
20 ctgaaagggc taaattgatt tccttaccaa ctagtaaaaa ctggaccttt ggaccccaa 2520
atatagatga actgatcttt atgttaagt acagccctgg ggtcatgtgc cggccctcac 2580
gagtcagaca gatgtttgct tccagagcct gtcggaagtc agtgatgatt ggaacggcgc 2640
tcaatgcgag cgagatgaag aagctcatca cccacatggg tgagatggac cacccttgga 2700
actgccccca cggcaggcca accatgaggg acgttgccaa tctggatgtc atctctcaga 2760
25 actgacacac ccctgttagc atagagttta ttacagattg ttcggtttgc aaagagaagg 2820
ttttaagtaa tctgattatc gttgtacaaa aattagcatg ctgctttaat gtactggatc 2880
catttaaaag cagtgttaag gcaggcatga tggagtgttc ctctagctca gctacttggg 2940
tgatccgggt ggagctcatg tgagcccagg actttgagac cactccgagc cacattcatg 3000
agactcaatt caaggacaaa aaaaaaaga tatttttgaa gccttttaaa aaaaaa 3056

30 human PMS2 protein

MKQLPAATVR LLSSSQIITS VVSVKELIE NSLDAGATSV DVKLENYGFD KIEVRDNGEG 60
IKAVDAPVMA MKYYTSKINS HEDLENLTTY GFRGEALGSI CCIAEVLITT RTAADNFSTQ 120
YVLDGSGHIL SQKPSHLGGG TTVTALRLFK NLPVRKQFYS TAKKCKDEIK KIQDLLMSFG 180
35 ILKPDRLIVF VHNKAVIWQK SRVSDHKMAL MSVLGTAVMN NMESFQYHSE ESQIYLSGFL 240
PKCDADHSFT SLSTPERSFI FINSRPVHQK DILKLIRHHY NLKCLKESTR LYPVFFLKID 300
VPTADVVDNL TPDKSQVLLQ NKESVLIALE NLMTTCYGPL PSTNSYENNK TDVSAADIVL 360
SKTAETDVLV NKVESGSKNY SNVDTSVIPF QNDMHNDESG KNTDDCLNHQ ISIGDFGYGH 420
CSSEISNIDK NTKNAFQDIS MSNVSWENSQ TEYSKTCFIS SVKHTQSENG NKDHIDESGE 480
40 NEEAEGLNS SEISADEWSR GNILKNSVGE NIEPVKILVP EKSLPCKVSN NNYPIPEQMN 540
LNEDSCNKKV NVIDNKSGKV TAYDLLSNRV IKKPMASAL FVQDHRPQFL IENPKTSLED 600
ATLQIEELWK TLSEEEKLKY EEKATKDLEK YNSQMKRAIE QESQMSLKDG RKKIKPTSAW 660
NLAQKHKLKT SLSNQPKLDE LLQSQIEKRR SQNIKMVQIP FSMKNLKINF KKQNKVDLEE 720
KDEPCLIHNL RFPDAWLMTS KTEVMLNLPY RVEEALLFKR LLENHKLPAE PLEKPIMLTE 780
45 SLFNGSHYLD VLYKMTADDQ RYSGSTYLSL PRLTANGFKI KLIPGVSITE NYLEIEGMAN 840
CLPFYGVADL KEILNAILNR NAKEVYECRP RKVISYLEGE AVRLSRQLPM YLSKEDIQDI 900
IYRMKHQFGN EIKECVHGRP FFHHLTYLPE TT 932

5 Human PMS2 cDNA
cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aaggccatca aacctattga tcggaagtca gtccatcaga tttgctctgg gcaggtggta 120
ctgagtctaa gcactgcggt aaaggagtta gtagaaaaca gtctggatgc tgggtccact 180
aatattgac taaagcttaa ggactatgga gtggatctta ttgaagttc agacaatgga 240
10 tgtgggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggct ttcgggggga agctctgagc 360
tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcatcggc gaaggttggg 420
actcgactga tgtttgatca caatgggaaa attatccaga aaacccccta ccccgcccc 480
agagggacca cagtcagcgt gcagcagtta tttccacac tacctgtgcg ccataaggaa 540
15 tttcaaagga atattaagaa ggagtatgcc aaaatgggtc aggtcttaca tgcatactgt 600
atcatttcag caggcatccg tgtaagttgc accaatcagc ttggacaagg aaacgcacag 660
cctgtgttat gcacaggtgg aagcccagc ataaaggaaa atatcggtc tgtgtttggg 720
cagaagcagt tgcaagcct cattcctttt gttcagctgc ccctagtga ctccgtgtgt 780
gaagagtacg gtttgagctg ttcggatgct ctgcataatc ttttttacct ctcaggtttc 840
20 atttcacaat gcacgcatgg agttggaagg agttcaacag acagacagtt tttctttatc 900
aaccggcgcc cttgtgacct agcaaaggct tgcagactcg tgaatgaggt ctaccacatg 960
tataatcgac accagtatcc atttggtgtt cttaacattt ctgttgatc agaatgcgtt 1020
gatatcaatg ttactccaga taaaaggcaa attttgctac aagaggaaaa gcttttgttg 1080
gcagttttaa agacctcttt gataggaatg tttgatagtg atgtcaacaa gctaaatgtc 1140
25 agtcagcagc cactgctgga tgttgaagg aacttaataa aaatgcatgc agcggatttg 1200
gaaaagccca tggtagaaaa gcaggatcaa tccccttcat taaggactgg agaagaaaaa 1260
aaagacgtgt ccatttccag actgcgagag gccttttctc ttcgtcacac aacagagAAC 1320
aagcctcaca gcccaaagac tccagaacca agaaggagcc ctctaggaca gaaagggggt 1380
atgctgtctt ctagcacttc aggtgccatc tctgacaaa gcgtcctgag acctcagaaa 1440
30 gaggcagtga gttccagtca cggaccagc gaccctacgg acagagcggg ggtggagaag 1500
gactcggggc acggcagcac ttccgtggat tctgaggggt tcagcatccc agacacgggc 1560
agtcactgca gcagcgagta tgcggccagc tcccagggg acaggggctc gcaggaaat 1620
gtggactctc aggagaaaag gcctgaaact gacgactctt tttcagatgt ggactgccat 1680
tcaaaccagg aagataccgg atgtaaattt cgagttttgc ctcagccaac taatctcgca 1740
35 accccaaaaca caaagcgttt taaaaaagaa gaaattcttt ccagttctga catttgtcaa 1800
aagttagtaa atactcagga catgtcagcc tctcagggtg atgtagctgt gaaaattaat 1860
aagaaagtgt tgcccctgga cttttctatg agttctttag ctaaacgaat aaagcagtta 1920
catcatgaag cacagcaaag tgaaggggaa cagaattaca ggaagttag ggcaaagatt 1980
tgtcctggag aaaatcaagc agccgaagat gaactaagaa aagagataag taaaacgatg 2040
40 tttgcagaaa tggaaatcat tggtcagttt aacctgggat ttataataac caaactgaat 2100
gaggatatct tcatagtgga ccagcatgcc acggacgaga agtataactt cgagatgctg 2160
cagcagcaca ccgtgtctca ggggcagagg ctcatagcac ctcagactct caacttaact 2220
gctgttaatg aagctgttct gatagaaaat ctggaaatat ttagaaagaa tggctttgat 2280
tttgttatcg atgaaaatgc tccagtcact gaaagggcta aactgatttc cttgccaact 2340
45 agtaaaaact ggaccttcgg accccaggac gtcgatgaac tgatcttcat gctgagcgac 2400
agccctgggg tcatgtgccg gccttcccga gtcgaagcaga tgtttgcctc cagagcctgc 2460
cggaagtcgg tgatgattgg gactgctctt aacacaagcg agatgaagaa actgatcacc 2520
cacatggggg agatggacca cccctggaac tgtcccatg gaaggccaac catgagacac 2580
atcgccaacc tgggtgtcat ttctcagAAC tgaccgtagt cactgtatgg aataattggt 2640

tttatcgag atttttatgt ttgaaagac agagtcttca ctaacctttt ttgttttaaa 2700
atgaaacctg ctacttaaaa aaaatacaca tcacacccat ttaaaagtga tcttgagaac 2760
cttttcaaac c
2771

5

human PMS1 protein

10 MKQLPAATVR LLSSSQIITS VVSVKELIE NSLDAGATSV DVKLENYGFD KIEVRDNGEG 60
IKAVDAPVMA MKYYTSKINS HEDLENLTTY GFRGEALGSI CCIAEVLITT RTAADNFSTQ 120
YVLDGSGHIL SQKPSHLGQG TTVTALRLFK NLPVRKQFYS TAKKCKDEIK KIQDLMSFG 180
ILKPDRLRIV VHNKAVIQK SRVSDHKMAL MSVLGTAVMN NMESFYHSE ESQIYLSGFL 240
PKCADHSFT SLSTPERSFI FINSRPVHOK DILKLIRHHY NLKCLKESTR LYPVFFLKID 300
VPTADVVDNL TPDKSQVLLQ NKESVLIALE NLMTCYGPL PSTNSYENNK TDVSAADIVL 360
15 SKTAETDVLV NKVESSGKNY SNVDTSVIPF QNDMHNDESG KNTDDCLNHQ ISIGDFGYGH 420
CSSEISNIDK NTKNAFQDIS MSNVSWENSQ TEYSKTCFIS SVKHTQSENG NKDHIDESGE 480
NEEEAGLENS SEISADEWSR GNILKNSVGE NIEPVKILVP EKSLPCKVSN NNYPIPEQMN 540
LNEDSCNKS NVIDNKGKV TAYDLLSNRV IKKPSASAL FVQDHRPQFL IENPKTSLED 600
ATLQIEELWK TLSEEEKLKY EEKATKDLER YNSQMKRAIE QESQMSLKDG RKKIKPTSAA 660
20 NLAQKHKLKT SLSNQPKLDE LLQSQIEKRR SQNIKMVQIP FSMKNLKINF KKQNKVDLEE 720
KDEPCLIHNL RFPDAWLMTS KTEVMLNPFY RVEEALLFKR LLENHKLPAE PLEKPIMLTE 780
SLFNGSHYLD VLYKMTADDQ RYSGSTYLSL PRLTANGFKI KLIPGVSITE NYLIEGEMAN 840
CLPFYGVADL KEIINAILNR NAKEVYECRP RKVISYLEGE AVRLSRQLPM YLSKEDIQDI 900
IYRMKHQFGN EIKECVHGRP FFHHLTYLPE TT
932

25

Human PMS1 cDNA

ggcagcagtg gctgcttgcg gctagtggtg ggtaattgcc tgcctcgcg tagcagcaag 60
ctgctctgtt aaaagcgaaa atgaacaat tgcctgcggc aacagttcga ctctttcaa 120
gttctcagat catcacttcg gtggtcagtg ttgtaaaaga gcttattgaa aactccttg 180
30 atgctggtgc cacaagcgta gatgttaaac tggagaacta tggatttgat aaaattgagg 240
tgcgagataa cggggagggt atcaaggctg ttgatgcacc tgtaatggca atgaagtact 300
acacctcaaa aataaatagt catgaagatc ttgaaaattt gacaacttac ggttttcgtg 360
gagaagcctt ggggtcaatt tgttgtatag ctgaggtttt aattacaaca agaacggctg 420
ctgataattt tagcaccacg tatgttttag atggcagtg ccacatactt tctcagaaac 480
35 cttcacatct tgggtcaagg acaactgtaa ctgctttaag attatttaag aatctactg 540
taagaaagca gttttactca actgcaaaaa aatgtaaaga tgaaataaaa aagatccaag 600
atctcctcat gagctttggt atccttaaac ctgacttaag gattgtcttt gtacataaca 660
aggcagttat ttggcagaaa agcagagtat cagatcaca gatggctctc atgtcagttc 720
tggggactgc tgttatgaac aatatggaat cctttcagta cactctgaa gaatctcaga 780
40 tttatctcag tggatttctt ccaaagtgtg atgcagacca ctctttcact agtctttcaa 840
caccagaaaag aagtttcac ttcataaaca gtcgaccagt acatcaaaaa gatattctaa 900
agttaaatccg acatcattac aatctgaaat gcctaaagga atctactcgt ttgtatcctg 960
ttttctttct gaaaatcgat gttctacag ctgatgttga tgtaaattta acaccagata 1020
aaagccaagt attattacaa aataaggaat ctgttttaat tgctcttgaa aatctgatga 1080
45 cgacttgtaa tggaccatta ctagtacaa attcttatga aaataataaa acagatgttt 1140
ccgcagctga catcgttctt agtaaaacag cagaaacaga tgtgcttttt aataaagtgg 1200
aatcatctgg aaagaattat tcaaatgttg atacttcagt cattccattc caaaatgata 1260
tgcataatga tgaatctgga aaaaacactg atgattgttt aaatcaccag ataagtattg 1320
gtgactttgg ttatgggtcat tgtagtagtg aaatttctaa cattgataaa aacactaaga 1380

atgcatttca ggacatttca atgagtaatg tatcatggga gaactctcag acggaatata 1440
gtaaaacttg ttttataagt tccgttaagc acaccagtc agaaaatggc aataaagacc 1500
atatagatga gagtggggaa aatgaggaag aagcaggtct tgaaaactct tcggaaattt 1560
ctgcagatga gtggagcagg ggaatatatac ttaaaaattc agtgggagag aatattgaac 1620
5 ctgtgaaaaa tttagtgcct gaaaaaagt taccatgtaa agtaagtaat aataattatc 1680
caatccctga acaaatgaat cttaatgaag attcatgtaa caaaaaatca aatgtaatag 1740
ataataaatc tggaaaagt acagcttatg atttacttag caatcgagta atcaagaaac 1800
ccatgtcagc aagtgcctct tttgttcaag atcatcgctc tcagtttctc atagaaaatc 1860
ctaagactag tttagaggat gcaacactac aaattgaaga actgtggaag acattgagtg 1920
10 aagaggaaaa actgaaatat gaagagaagg ctactaaaga cttggaacga tacaatagtc 1980
aaatgaagag agccattgaa caggagtcac aaatgtcact aaaagatggc agaaaaaaga 2040
taaaacccac cagcgcatgg aatttggccc agaagcaca gttaaaaacc tcattatcta 2100
atcaacaaaa acttgatgaa ctcttccagt cccaaattga aaaaagaagg agtcaaaata 2160
ttaaataatg acagatcccc ttttctatga aaaacttaaa aataaatttt aagaacaaaa 2220
15 acaaagtga cttagaagag aaggatgaac cttgcttgat ccacaatctc aggtttcctg 2280
atgcatggct aatgacatcc aaaacagagg taatgttatt aaatccatat agagtagaag 2340
aagccctgct atttaaaaga cttcttgaga atcataaact tcctgcagag ccactggaaa 2400
agccaattat gttaacagag agtcttttta atggatctca ttatttagac gttttatata 2460
aaatgacagc agatgaccaa agatacagtg gatcaactta cctgtctgat cctcgtctta 2520
20 cagcgaatgg tttcaagata aaattgatac caggagtctc aattactgaa aattacttgg 2580
aaatagaagg aatggctaata tgtctcccat tctatggagt agcagattta aaagaaattc 2640
ttaatgctat attaaacaga aatgcaaagg aagtttatga atgtagacct cgcaaagtga 2700
taagttatct agagggagaa gcagtgcgtc tatccagaca attacccatg tacttatcaa 2760
aagaggacat ccaagacatt atctacagaa tgaagcacca gtttggaat gaaattaaag 2820
25 agtgtgttca tggctgcccc ttttttcatc atttaaccta tcttccagaa actacatgat 2880
taaatatggt taagaagatt agttaccatt gaaattgggt ctgtcataaa acagcatgag 2940
tctgtgttta aattatcttt gtattatgtg tcacatgggt attttttaaa tgaggattca 3000
ctgacttggt tttatattga aaaaagtcc acgtattgta gaaaacgtaa ataaactaat 3060
aac 3063
30

human MSH2 protein
MAVQPKETLQ LESAAEVGFV RFFQGMPEKP TTTVRLFDRG DFYTAHGEDA LLAAREVFKT 60
QGVIKYMGPA GAKNLQSVVL SKMNFESFVK DLLLRQYRV EVYKNRAGNK ASKENDWYLA 120
35 YKASPGNLSQ FEDILFGNND MSASIGVVG V KMSAVDGRQ VGVGYVDSIQ RKLGLCEFPD 180
NDQFSNLEAL LIQIGPKECV LPGAETAGDM GKLRQIIQRG GILITERKKA DFSTKDIYQD 240
LNRLKGGKG EQMNSAVLPE MENQVAVSSL SAVIKFLELL SDDSNGQFE LTTFDFSQYM 300
KLDIAAVRAL NLFQGSVEDT TGSQSLAALL NKCKTPQGQR LVNQWIKQPL MDKNRIEERL 360
NLVEAFVEDA ELRQLQEDL LRRFPDLNRL AKKFQRQAAN LQDCYRLYQG INQLPNVIQA 420
40 LEKHGKHQK LLLAVFVTPL TDLRSDFSKF QEMIETTLDM DQVENHEFLV KPSFDPNLSE 480
LREIMNDLEK KMQSTLISAA RDLGLDPGKQ IKLDSSAQFG YYFRVTCKEE KVLNNKNFS 540
TVDIQKNGVK FTNSKLTSLN EBYTKNKEY EEAQDAIVKE IVNISSGYVE PMQTLNDVLA 600
QLDAVVSFAH VSNAPVPYV RPAILEKGQ RIIKASRHA CVEVQDEIAF IPNDVYFEKD 660
KQMFHIIITGP NMGGKSTYIR QTVIVLMAQ IGCVPCESA EVSIVDCILA RVGAGDSQLK 720
45 GVSTFMAEML ETASILRSAT KDSLIIDEL GRGTSTYDGF GLAWAISEYI ATKIGAFCMF 780
ATHFHELTAL ANQIPTVNNL HVTALTTEET LTMLYQVKKG VCDQSFGIHV AELANFPKHV 840
IECAKQKALE LEEFYIGES QGYDIMEPAA KKCYLEREQG EKIIQEFLSK VKQMPFTEMS 900
EENITIKLKQ LKAEVIKNN SFVNEIISRI KVT 934

Human MSH2 cDNA

ggcgggaaac agcttagtgg gtgtgggggc gcgcattttc ttcaaccagg aggtgaggag 60
 gtttcgacat ggcggtgcag ccgaaggaga cgctgcagtt ggagagcgcg gccgaggtcg 120
 gcttcgtgcg cttctttcag ggcatgccgg agaagccgac caccacagtg cgccttttcg 180
 5 accggggcga cttctatacg gcgcacggcg aggacgcgct gctggccgcc cgggaggtgt 240
 tcaagaccca gggggtgac aagtacatgg ggccggcagg agcaaagaat ctgcagagtgt 300
 ttgtgcttag taaaatgaat tttgaatctt ttgtaaaaga tcttctctctg gttcgtcagt 360
 atagagtga agtttataag aatagagctg gaaataaggc atccaaggag aatgatttgt 420
 atttggcata taaggcttct cctggcaatc tctctcagtt tgaagacatt ctctttggta 480
 10 acaatgatat gtcagcttcc attggtgttg tgggtgttaa aatgtccgca gttgatggcc 540
 agagacaggt tggagttggg tatgtggatt ccatacagag gaaactagga ctgtgtgaat 600
 tccttgataa tgatcagttc tccaatcttg aggtctctct catccagatt ggaccaaagg 660
 aatgtgtttt acccgaggga gagactgctg gagacatggg gaaactgaga cagataattc 720
 aaagaggagg aattctgac acagaaagaa aaaaagctga cttttccaca aaagacattt 780
 15 atcaggacct caaccggttg ttgaaaggca aaaagggaga gcagatgaat agtgctgtat 840
 tgcagaaat ggagaatcag gttgcagttt catcactgtc tgcgtaatc aagtttttag 900
 aactcttacc agatgattcc aactttggac agtttgaact gactactttt gacttcagcc 960
 agtatatgaa attggatatt gcagcagtc gagcccttaa cctttttcag ggttctgttg 1020
 aagataccac tggctctcag tctctggctg ccttgctgaa taagtgtaaa acccctcaag 1080
 20 gacaaagact tgtaaccag tggattaagc agcctctcat ggataagaac agaataagagg 1140
 agagattgaa tttagtggaa gctttttag aagatgcaga attgaggcag actttacaag 1200
 aagatttact tcgtcgattc ccagatctta accgacttgc caagaagttt caaagacaag 1260
 cagcaaaactt acaagattgt taccgactct atcaggggat aaatcaacta cctaattgta 1320
 tacaggctct ggaaaaacat gaaggaaaac accagaaatt attgttggca gtttttgtga 1380
 25 ctctcttacc tgatcttctg tctgacttct ccaagtttca ggaaatgata gaaacaactt 1440
 tagatatgga tcagggtgaa aaccatgaat tccttgtaaa accttcattt gatcctaata 1500
 tcagtgaatt aagagaaata atgaatgact tggaaaagaa gatgcagtc acattaataa 1560
 gtgcagccag agatcttggc ttggaccctg gcaaacagat taaactggat tccagtgcac 1620
 agtttgata ttaacttctg gtaacctgta aggaagaaaa agtccttctg aacaataaaa 1680
 30 actttagtac tgtagatata cagaagaatg gtgttaaat taccacagc aaattgactt 1740
 ctttaaatga agagtatacc aaaaataaaa cagaatatga agaagcccag gatgccattg 1800
 ttaagaaat tgtcaatatt tcttcaggct atgtagaacc aatgcagaca ctcaatgatg 1860
 tgttagctca gctagatgct gttgtcagct ttgctcacgt gtcaaatgga gcacctgttc 1920
 catatgtacg accagccatt ttggagaaag gacaaggag aattatatta aaagcatcca 1980
 35 ggcattgctt gtttgaagtt caagatgaaa ttgcatttat tcctaatagac gtatactttg 2040
 aaaaagataa acagatgttc cacatcatta ctggcccaa tatgggaggt aaatcaacat 2100
 atattcgaca aactggggtg atagtactca tggcccaaat tgggtgtttt gtgccatgtg 2160
 agtcagcaga agtgtccatt gtggactgca tcttagcccg agtaggggct ggtgacagtc 2220
 aattgaaagg agtcctcacg ttcatggctg aaatgttga aactgcttct atcctcaggt 2280
 40 ctgcaaccaa agattcatta ataatcatag atgaattggg aagaggaact tctacctacg 2340
 atggatttgg gttagcatgg gctatatcag aatacatgac aacaagatt ggtgcttttt 2400
 gcatgtttgc aaccatttt catgaactta ctgccttggc caatcagata ccaactgtta 2460
 ataacttaca tgtcacagca ctccacctg aagagacctt aactatgctt tatcagggtga 2520
 agaaagggtg ctgtgatcaa agttttggga ttcatgttgc agagcttgct aatttcccta 2580
 45 gacatgtaat agagtgtgct aaacagaaag ccctggaact tgaggagttt cagtatattg 2640
 agaaatcgca aggatatgat atcatggaac cagcagcaaa gaagtgttat ctggaagag 2700
 agcaaggtga aaaaattatt caggagttcc tgtccaaggt gaaacaaatg ccctttactg 2760
 aaatgtcaga agaaaacatc acaataaagt taaaacagct aaaagctgaa gtaatagcaa 2820
 agaataatag ctttgtaaat gaaatcattt cacgaataaa agttactacg tgaaaaatcc 2880

5 cagtaatgga atgaaggtaa tattgataag ctattgtctg taatagtttt atattgtttt 2940
 atattaaccc tttttccata gtgttaactg tcagtgccca tgggctatca acttaataag 3000
 atatttagta atattttact ttgaggacat tttcaaagat ttttattttg aaaaatgaga 3060
 gctgtaactg aggactgttt gcaattgaca taggcaataa taagtgatgt gctgaatttt 3120
 ataaataaaa tcatgtagtt tgtgg
 3145

human MLH1 protein
 10 MSFVAGVIRR LDETUVNRIA AGEVIQRPAN AIKEMIENCL DAKSTSIQVI VKEGGLKLIQ 60
 IQDNGTGIRK EDLDIVCERF TTSKLQSFED LASISTYGFR GEALASISHV AHVTITTKTA 120
 DGKCAIRASY SDGKLKAPPK PCAGNQGTQI TVEDLFYNIA TRRKALKNPS EEYGKILEVV 180
 GRYSVHNAGI SFSVKKQGET VADVRTL PNA STVDNIRSIF GNAVSRELIE IGCEDKTLAF 240
 KMNGYISNAN YSVKCCIFLL FINHRLVEST SLRKAIVTVY AAYLPKNTHP FLYLSLEISP 300
 QNVDEVNVHPT KHEVHFLHEE SILERVQQHI ESKLLGSNSS RMYFTQTLLP GLAGPSGEMV 360
 15 KSTTSLTSSS TSGSSDKVYA HQMVRTDSRE QKLD AFLQPL SKPLSSQPQA IVTEDKTDIS 420
 SGRARQDEE MLELPAPAEV AAKNQSLEGD TTKGTSEMSE KRGPTSSNPR KHRHRESDVE 480
 MVEDDSRKEM TAACTPRRRI INLTSVLSLQ EEINEQGHEV LREMLHNHSF VGCVPQWAL 540
 AQHQTKLYLL NTKLSEELF YQILIYDFAN FGVRLSEPA PLFDLAMLAL DSPESGWTEE 600
 DGPKEGLAEY IVEFLKKKAE MLADYFSLEI DEEGNLIGLP LLIDNYVPPL EGLPIFILRL 660
 20 ATEVNWDEEK ECFESLSKEC AMFY SIKRQY ISEESTLSGQ QSEVPGSIPN SWKWTVEHIV 720
 YKALRSHILP PKHFTEDGNI LQLANLPDLY KVFERC 756

Human MLH1 cDNA
 25 cttggctcct ctggcgccaa aatgctgttc gtggcagggg ttattcggcg gctggacgag 60
 acagtggtga accgcatcgc ggcgggggaa gttatccagc ggccagctaa tgctatcaaa 120
 gagatgattg agaactgttt agatgcaaaa tccacaagta ttcaagtgat tgtaaaagag 180
 ggaggcctga agttgattca gatccaagac aatggcaccg ggatcaggaa agaagatctg 240
 gatattgtat gtgaaagggt cactactagt aaactgcagt cctttgagga tttagccagt 300
 30 atttctacct atggctttcg aggtgaggct ttggccagca taagccatgt ggctcatgtt 360
 actattacaa cgaaaacagc tgatggaaaag tgtgcataca gagcaagtta ctcagatgga 420
 aaactgaaag cccctcctaa accatgtgct ggcaatcaag ggaccagat cacggtggag 480
 gaccctttttt acaacatagc cacgaggaga aaagctttaa aaaatccaag tgaagaatat 540
 gggaataattt tggaagtgtt tggcagggtat tcagtacaca atgcaggcat tagtttctca 600
 gttaaaaaac aaggagagac agtagctgat gttaggacac taccatagc ctcaaccgtg 660
 35 gacaatatct gctccatctt tggaatgctt gttagtcgag aactgataga aattggatgt 720
 gaggataaaa ccctagcctt caaaatgaat ggttacatat ccaatgcaaa ctactcagtg 780
 aagaagtgca tcttcttact cttcatcaac catcgtctgg tagaatcaac ttccttgaga 840
 aaagccatag aaacagtgtg tgcagcctat ttgccccaaa acacacaccc attcctgtac 900
 ctcagtttag aaatcagtc ccagaatgtg gatgttaatg tgcacccac aaagcatgaa 960
 40 gttcacttcc tgcacgagga gagcatcctg gagcgggtgc agcagcacat cgagagcaag 1020
 ctccctgggt ccaattcctc caggatgtac ttcaccaga ctttgctacc aggactgtct 1080
 ggcccctctg gggagatggt taaatccaca acaagtctga cctcgtcttc tacttctgga 1140
 agtagtgata aggtctatgc ccaccagatg gttcgtacag attcccggga acagaagctt 1200
 gatgcatttc tgcagcctct gagcaaaacc ctgtccagtc agcccaggc cattgtcaca 1260
 45 gaggataaga cagatatctt tagtggcagg gctaggcagc aagatgagga gatgcttgaa 1320
 ctcccagccc ctgctgaagt ggctgccaaa aatcagagct tggaggggga tacaacaaag 1380
 gggacttcag aaatgtcaga gaagagagga cctacttoca gcaacccag aaagagacat 1440
 cggaagatt ctgatgtgga aatggtggaa gatgattccc gaaaggaaat gactgcagct 1500
 tgtaccccc ggagaaggat cattaacctc actagtgttt tgagtctcca ggaagaaatt 1560

aatgagcagg gacatgaggt tctccgggag atgttgcata accactcctt cgtgggctgt 1620
gtgaatcctc agtgggcctt ggcacagcat caaaccaagt tataccttct caacaccacc 1680
aagcttagtg aagaactgtt ctaccagata ctacattatg attttgcaa ttttggtgtt 1740
ctcaggttat cggagccagc accgctcttt gaccttgcca tgcttgctt agatagtcca 1800
5 gagagtggct ggacagagga agatggctcc aaagaaggac ttgctgaata cattgttgag 1860
tttctgaaga agaaggctga gatgcttgca gactatttct ctttggaat tgatgaggaa 1920
gggaacctga ttggattacc ccttctgatt gacaactatg tgccccctt ggagggactg 1980
cctatcttca ttcttcgact agccactgag gtgaattggg acgaagaaaa ggaatgttt 2040
gaaagcctca gtaaagaatg cgctatgttc tattccatcc ggaagcagta catatctgag 2100
10 gactcgaccc tctcaggcca gcagagtga gtgcctggct ccattccaaa ctctggaag 2160
tggtgtgtg aacacattgt ctataaagcc ttgcgctcac acattctgcc tcctaaacat 2220
ttcacagaag atggaaatat cctgcagctt gctaacctgc ctgactata caaagtctt 2280
gagagtggtt aaatatggtt atttatgcac tgtgggatgt gttcttctt ctctgtattc 2340
cgatacaaa tggtgtatca aagtgtgata taaaaagtgt accaacataa gtgttggtag 2400
15 cacttaagac ttatacttgc cttctgatag tattccttta tacacagtgg attgattata 2460
aataaataga tgtgtcttaa cata 2484

hPMS2-134 protein
20 MKQLPAATVR LLSSSQIITS VVSVVKELIE NSLDAGATSV DVKLENYGFD KIEVRDNNEG 60
IKAVDAPVMA MKYYTSKINS HEDLENLTYY GFRGEALGSI CCIAEVLITT RTAADNFSTQ 120
YVLDGSGHIL SQK 133

hPMS2-134 cDNA
25 cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aagccatca aacctattga tcggaagtca gtccatcaga tttgctctgg gcaggtggtg 120
ctgagtctaa gcactgcggt aaaggagttg gtagaaaaca gtctggatgc tgggtgccact 180
aatattgatc taaagcttaa ggactatgga gtggatctta ttgaagtctc agacaatgga 240
tgtggggtag aagaagaaaa ctctgaagcc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggtt ttcgggggga agctctgagc 360
30 tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcatcgcc gaaggttgga 420
acttga 426

hMSH6 (human cDNA) ACCESSION U28946
35 MSRQSTLYSFFPKSPALSDANKASARASREGGAAAAAGASPS
GGDAWSEAGPGFRPLARSASPPKAKNLNGLRRSVAPAPTSCDFSPGDLVWAKMEG
40 YPWWPCLVYNHPFDGTFIREKGKSVRVHVQFFDDSPTRGWVSKRLKPYTGSKSKEAQ
KGGHFYSAPKEILRAMQRADEALNKDKIKRLELAVCDEPSEPEEEEEMEVGTTYVTDK
SEEDNEIESEEEVQPKTQGSRRSRQIKRRVISDSSESDIGGSDVEFKPDTKEEGSSD
45 EISSGVGDSESEGLNSPVKVARKRKRMVTGNGSLKRKSRKETPSATKQATSISETK
NTLRAFSAPQNSQAHVSGGDDSSRPTVWYHETLEWLKEEKRRDEHRRRDPDHPDFD
50 ASTLYVPEDFLNSCTPGMRKWWQIKSQNFDLVICYKVGKFEYELHMDALIGVSELGLV
FMKGNWAHSGFPEIAFGRYSDSLVQKGYKVARVEQTETPEMMEARCRKMAHISKYDRV
VRREICRIITKGTQTYSVLEGDPSENYSKYLLSLKEEEDSSGHTRAYGVCVDTSLG
55 KFFIGQFSDDRHCSTRFTLVAHYPPVQVLFKGNLSKETKITLSSLSLQEGELIPG
SQFDWASKTLRTLLLEEEYFREKLSDGIGVMLPQVLKGMTSESDSIGLTPGEKSELALS
60 ALGGCVFYLLKCLIDQELLSMANFEEYIPLSDTVSTTRSGAIFTKAYQRMVLDAVTL
NNLEIFLNGTNGSTEGTLLEVRVDTCHTPFGKRLKQWLCAPLCNHYAINDRLDAIEDL

5 MVVPDKISEVVELLKKLPDLERLLSKIHNVGSPLKSNHPDSRAIMYEETTSKKKII
 DFLSALEGFKVMCKIIIGIMEEVADGFKSKILKQVISLQTKNPEGRFPDLTVELNRWDT
 AFDHEKARKTGLITPKAGFSDYDQALADIRENEQSLEYLEKQRNRIGCRTIVYWGI
 10 GRNRYQLEIPENFTTRNLPPEEYELKSTKKGCKRYWTKTIEKKLANLINAERRDVSLK
 DCMRRLFYNFDDKNYKDWQSAVECIAVLVDVLLCLANYSRGGDGPMPRPVILLPEDTPPF
 LELKGSRHPICITKTFFGDDFIENDILIGCEEEQENGKAYCVLVTGPNMGGKSTLMRQ
 15 AGLLAVMAQMGCYVPAEVCRLTPIDRVFTRLGASDRIMSGESTFFVELSETASILMHA
 TAHSVLVDELGRGTATFDGTAIANAVVKELAETIKCRTLFSTHYHSLVEDYSQNVAV
 RLGHMACMVENECEDPSETITFLYKFIKGACPKSYGFNAARLANLPEEVIQKGRKA
 20 REFEBMNQSLRLFREVCLASERSTVDAEAVHKLLTLIKEL"

hPMSR2 (human cDNA) ACCESSION U38964

25 1 ggcgctccta cctgcaatg gctagtgcc agtgctgggc cgccgctect gccgtgcatg
 61 ttggggagcc agtacatgca ggtgggctcc acacggagag gggcgagac ccggtgacag
 121 ggctttacct ggtacatcgg catggcgcaa ccaagcaag agagggtggc gcgtgccaga
 181 caccacacgt cggaaccgac cagacaccaa cggtcgaaa ccgccaagac accaacgctc
 241 ggaaaccgac agacaccaac gctcgaaaac cgccagacac caaggctcgg aatccacgcc
 301 agggccacgac ggagggcgac tacctccctt ctgacctgct tgctggcgtt cggaaaaaac
 361 gcagtccggt gtgctctgat tggctccaggc tctttgacgt caccgactcg acccttgaca
 421 gagccactag gcgaaaagga gagacgggaa gtattttttc cgccccgccc ggaaggggtg
 481 gagcacaacg tcgaaagcag ccgttgggag ccaggaggc gggggcgctg tgggagccgt
 541 ggagggaact ttccagtcgc ccgaggcgga tccggtgttg catccttgga gcgagctgag
 601 aactcgagta cagaacctgc taaggccatc aaacctattg atcggaagtc agtccatcag
 661 atttgctctg ggcgggtggt accgagtcta agccgaatg ccgtgaagga gttagtagaa
 721 aacagtctgg atgctggtgc cactaatgtt gatctaaagc ttaaggacta tggagtggat
 781 ctcatgtgaag ttccaggcaa tggatgtggg gtagaagaag aaaacttcga aggccttact
 841 ctgaaacatc acacatgtaa gattcaagag ttgcccagc taactcagg ggaacctttt
 901 ggcttttcgg gggaagctct gagctcactt tgtgactga gtgatgtcac catttctacc
 961 tgccgtgtat cagcgaaggt tgggactcga ctggtgtttg atcactatgg gaaaatcatc
 1021 cagaaaaacc cctacccccg ccccagagg atgacagtca gcgtgaagca gttattttct
 1081 acgctacctg tgcaccataa agaatttcaa aggaatatta agaagaaacg tgcctgcttc
 1141 cctcttcgct tctgcccgtg ttgtcagttt cctgaggcct ccccagccat gcttctgtg
 1201 cagcctgtag aactgactcc tagaagtacc ccacccacc cctgctcctt ggaggacaac
 1261 gtgatcactg tattcagctc tgtcaagaat ggtccaggtt cttctagatg atctgcacaa
 1321 atgggttcct tcctccttcc tgatgtctgc cattagcatt ggaataaagt tctgctgaa
 1381 aatccaaaaa aaaaaaaaaa aaaaaaaa

hPMSR2 (human protein) ACCESSION U38964

MAQPKQERVARARHQRSETARHQRSETAKTPTLGNRQTPTLGNR

55 QTPRLGIHARPRRRATSLTLLTLLAFGKNAVCALIGPGSLTSRTRPLTEPLGEKERR
 EVFFPPRPERVEHNVESSRWEPRRRGACGSRGNFSPRGGSGVASLERAENSSTEPA
 60 KAIKPIDRKSVMHQCISGVPVPSLRPNVAVKELVENSIDAGATNVDLKLKDYGVDLIEVS
 GNGCGVEEENFEGFTLKHHTCKIQEFADLTQVETFGFRGEALSSLCALSDVTISTCRV
 SAKVGTSLVFDHYGKIIQKTPYPRPRGMTVSVKQLFSTLPVHHKEFQRNIKKKRACFP
 65 FAFCRDCQFPEASPAMLPVQPVELTPRSTPPHPCSLEDNVITVFSSVKNGPGSSR

HPMSR3 (human cDNA) ACCESSION U38979

70 1 tttttagaaa ctgatgttta ttttccatca accatttttc catgctgctt aagagaatat
 61 gcaagaacag cttaagacca gtcagtgggt gctcctaccc attcagtggc ctgagcagtg
 121 gggagctgca gaccagtctt ccgtggcagg ctgagcgctc cagtcttcag tagggaattg
 181 ctgaataggc acagagggca cctgtacacc ttcagaccag tctgcaacct caggctgagt
 241 agcagtgaac tcaggagcgg gagcagtcca ttcacctga aattcctcct tggctactgc
 301 cttctcagca gcagcctgct cttctttttc aatctcttca ggatctctgt agaagtacag
 361 atcaggcatg acctcccatg ggtgttcacg ggaaatgggt ccacgcgatg gcagaacttc
 421 ccgagccagc atccaccaca ttaaacccac tgagtggagt ccctgtgtgt tgcaggggat
 481 ggcaatgtcc acatagcgca gaggagaatc tgtgttacac agcgcaatgg taggttaggtt

5 541 aacataagat gcctccgtga gagcggaagg ggcggcgga cccgggcctg gcccgatatgt
601 gtccttgagg gcctagacta ggccgtcgct gtatggtgag cccagggag gcggatctgg
661 gccccagaa ggacacccgc ctggatttgc cccgtagccc ggcccgggcc cctcgggagc
721 agaacagcct tggtaggtg gacaggagg gacctcgca gcagacgagc gcgcccagca
781 cagcagcccc gccccggcct ctggggagcc ggggggcaga ggctgcggag cccagggagg
841 gtctatcagc cacagtctct gcatgtttcc aagagcaaca ggaaatgaac acattgcagg
901 ggccagtgtc attcaaatgatt gtggctgtgg atttcaccca ggaggagtgg cggcaactgg
961 accctgatga gaagatagca tacggggatg tgatgttga gaactacagc catctagtgt
10 1021 ctgtggggtg tgattatcac caagccaaac atcatcatgg agtggagggtg aaggaaagtgg
1081 agcagggaga ggagccgtgg ataattggaag gtgaatttcc atgtcaacat agtccagaac
1141 ctgctaaggc catcaaacct attgatcgga agtcagttca tcagatttgc tctgggccag
1201 tggtagctag tctaagcact gcagtgaagg agttagtaga aaacagtctg gatgctggtg
1261 ccactaatat tgatctaaag cttaaggact atggagtga tctcattgaa gtttcagaca
1321 atggatgtgg ggtagaagaa gaaaactttg aaggcttaac ctctttcagc tctgaaacat
1381 cacacatgta agattcaaga gtttccgac ctaactgaag ttgaaacttt cggttttcag
1441 ggggaagctc tgagctcact gtgtgcaactg agcagtgta ccatttctac ctgcccagcg
15 1501 ttggtgaagg ttgggactcg actggtgttt gatcacgatg ggaaaatcat ccaggaaacc
1561 ccctaccccc accccagagg gaccacagtc agcgtgaagc agttattttc tacgctacat
1621 gtgcggcata aggaatttca aaggaatatt aagaagacgt gctgtcttcc ccttcgcctt
20 1681 ctgcgctgat tgcagtttc ctgaggcctc cccagccatg cttcctgtac agcctgcaga
1741 actgtgagtc aattaaacct cttttcttca taaattaaaa aaaaa

25 hPMSR3 (human protein) ACCESSION U38979

MCPWRPRLGRRCMVSPREADLGPQKDRDLPRSPARAPREQNS
30 LGEVDRRGPREQTRAPATAAPRPLGSRGAEAAEPQEGLSATVSACFQEQEMNTLQG
PVSFKDVAVDFTQEEWRQLDPDEKIAYGDMLENYSHLVSVGYDYHQAHHHGVEVKE
VEQGEPEWIMEGEFPCQHSPEPAKAIKPIDRKSVMHICSGPVVLSLSTAVKELVENS
35 DAGATNIDLKLDYGVDLIEVSDNGCGVEEENFEGLISFSSETSHM"

hPMSL9 (human cDNA) ACCESSION NM_005395

40 1 atgtgtcctt ggcggcctag actaggccgt cgctgtatgg tgagccccag ggaggcggat
61 ctggggcccc agaaggacac ccgcctggat ttgccccgta gcccgcccg ggccccctcg
121 gagcagaaca gccttggtga ggtggacagg aggggacctc gcgagcagac gcgcccagca
181 gcgacagcag ccccgccccg gcctctcggg agccgggggg cagaggctgc ggagccccag
241 gagggtctat cagccacagt ctctgcatgt ttccaagagc aacaggaaat gaacacattg
45 301 cagggggccag tgtcattcaa agatgtggct gtggatttca cccaggagga gtggcggcaa
361 ctggaccctg atgagaagat agcatacggg gatgtgatgt tggagaacta cagccatcta
421 gtttctgtgg ggtatgatta tcaccaagcc aaacatcatc atggagtga ggtgaaggaa
481 gtggagcagg gagaggagcc gtggataatg gaaggtgaat ttccatgtca acatagttca
541 gaacctgcta aggccatcaa acctattgat cggaagtcag tccatcagat ttgctctggg
50 601 ccagtgttac tgagtctaag cactgcagtg aaggagttag tagaaaaacag tctggatgct
661 ggtgccacta atattgatct aaagcttaag gactatggag tggatctcat tgaagtttca
721 gacaatggat gtggggtaga agaagaaaac tttgaaggct taatctcttt cagctctgaa
781 acatcacaca tgtaa

55 hPMSL9 (human protein) ACCESSION NM_005395

MCPWRPRLGRRCMVSPREADLGPQKDRDLPRSPARAPREQNS
60 LGEVDRRGPREQTRAPATAAPRPLGSRGAEAAEPQEGLSATVSACFQEQEMNTLQG
PVSFKDVAVDFTQEEWRQLDPDEKIAYGDMLENYSHLVSVGYDYHQAHHHGVEVKE
VEQGEPEWIMEGEFPCQHSPEPAKAIKPIDRKSVMHICSGPVVLSLSTAVKELVENS
65 DAGATNIDLKLDYGVDLIEVSDNGCGVEEENFEGLISFSSETSHM"

References

1. Attfield, P.V. and Pinney, R.J. 1985. Elimination of multicopy plasmid R6K by bleomycin. *Antimicrob. Agents Chemother.* 27(6):985-988.

5

2. Baker S.M., Bronner, C.E., Zhang, L., Plug, A.W., Robatez, M., Warren, G., Elliott, E.A., Yu, J., Ashley, T., Arnheim, N., Bradley, N., Flavell, R.A., and Liskay, R.M. 1995. Male defective in the DNA mismatch repair gene PMS2 exhibit abnormal chromosome synapsis in meiosis. *Cell* 82:309-319.

10

3. Berry, A. 1996. Improving production of aromatic compounds in *Escherichia coli* by metabolic engineering. *Trends Biotechnol.* 14(7):250-256.

15

4. Bronner, C.E., Baker, S.M., Morrison, P.T., Warren, G., Smith, L.G., Lescoe, M.K., Kane, M., Earabino, C., Lipford, J., Lindblom, A., Tannergard, P., Bollag, R.J., Godwin, A., R., Ward, D.C., Nordenskjold, M., Fishel, R., Kolodner, R., and Liskay, R.M. 1994. Mutation in the DNA mismatch repair gene homologue *hMLH1* is associated with hereditary non-polyposis colon cancer. *Nature* 368:258-261.

20

5. de Wind N., Dekker, M., Berns, A., Radman, M., and Riele, H.T. 1995. Inactivation of the mouse *Msh2* gene results in mismatch repair deficiency, methylation tolerance, hyperrecombination, and predisposition to cancer. *Cell* 82:321-300.

25

6. Drummond, J.T., Li, G.M., Longley, M.J., and Modrich, P. 1995. Isolation of an hMSH2-p160 heterodimer that restores mismatch repair to tumor cells. *Science* 268:1909-1912.

30

7. Drummond, J.T., Anthoney, A., Brown, R., and Modrich, P. 1996. Cisplatin and adriamycin resistance are associated with MutLa and mismatch repair deficiency in an ovarian tumor cell line. *J.Biol.Chem.* **271**:9645-19648.
8. Edelmann, W., Cohen, P.E., Kane, M., Lau, K., Morrow, B., Bennett, S., Umar, A., Kunkel, T., Cattoretti, G., Chagnatti, R., Pollard, J.W., Kolodner, R.D., and Kucherlapati, R. 1996. Meiotic pachytene arrest in MLH1-deficient mice. *Cell* **85**:1125-1134.
9. Fishel, R., Lescoe, M., Rao, M.R.S., Copeland, N.J., Jenkins, N.A., Garber, J., Kane, M., and Kolodner, R. 1993. The human mutator gene homolog *MSH2* and its association with hereditary nonpolyposis colon cancer. *Cell* **7**:1027-1038.
10. Fu, K.P., Grace, M.E., Hsiao, C.L., and Hung, P.P. 1988. Elinination of antibiotic-resistant plasmids by quinolone antibiotics. *Chemotherapy* **34**(5):415-418.
11. Leach, F.S., Nicolaides, N.C, Papadopoulos, N., Liu, B., Jen, J., Parsons, R., Peltomaki, P., Sistonen, P., Aaltonen, L.A., Nystrom-Lahti, M., Guan, X.Y., Zhang, J., Meltzer, P.S., Yu, J.W., Kao, F.T., Chen, D.J., Cerosaletti, K.M., Fournier, R.E.K., Todd, S., Lewis, T., Leach R.J., Naylor, S.L., Weissenbach, J., Mecklin, J.P., Jarvinen, J.A., Petersen, G.M., Hamilton, S.R., Green, J., Jass, J., Watson, P., Lynch, H.T., Trent, J.M., de la Chapelle, A., Kinzler, K.W., and Vogelstein, B. 1993. Mutations of a *mutS* homolog in hereditary non-polyposis colorectal cancer. *Cell* **75**:1215-1225.

12. Li, G.-M., and Modrich, P. 1995. Restoration of mismatch repair to nuclear extracts of H6 colorectal tumor cells by a heterodimer of human mutL homologs. *Proc. Natl. Acad. Sci. USA* **92**:1950-1954.
- 5
13. Liu, B., Parsons, R., Papadopoulos, N., Nicolaides, N.C., Lynch, H.T., Watson, P., Jass, J.R., Dunlop, M., Wyllie, A., Peltomaki, P., de la Chapelle, A., Hamilton, S.R., Vogelstein, B., and Kinzler, K.W. 1996. Analysis of mismatch repair genes in hereditary non-polyposis colorectal cancer patients. *Nat. Med.* **2**:169-174.
- 10
14. Modrich, P. 1994. Mismatch repair, genetic stability, and cancer. *Science* **266**:1959-1960.
- 15
15. Nicolaides, N.C., Gualdi, R., Casadevall, C., Manzella, L., and Calabretta, B. 1991. Positive autoregulation of *c-myb* expression via MYB binding in the 5' flanking region of the human *c-myb* gene. *Mol. Cell. Biol.* **11**:6166-6176.
- 20
16. Nicolaides, N.C., Correa, I., Casadevall, C., Travali, S., Soprano, K.J., and Calabretta, B. 1992. The Jun family members, c-JUN and JUND, transactivate the human *c-myb* promoter via an Ap1 like element. *J. Biol. Chem.* **267**, 19665-19672.
- 25
17. Nicolaides, N.C., Papadopoulos, N., Liu, B., Wei, Y.F., Carter, K.C., Ruben, S.M., Rosen, C.A., Haseltine, W.A., Fleischmann, R.D., Fraser, C.M., Adams, M.D., Venter, C.J., Dunlop, M.G., Hamilton, S.R., Petersen, G.M., de la Chapelle, A., Vogelstein, B., and kinzler, K.W. 1994. Mutations of two PMS homologs in hereditary nonpolyposis colon cancer. *Nature* **371**: 75-80.
- 30

18. Nicolaides N.C., Kinzler, K.W., and Vogelstein, B. 1995. Analysis of the 5' region of PMS2 reveals heterogenous transcripts and a novel overlapping gene. *Genomics* **29**:329-334.

5

19. Nicolaides, N.C., Carter, K.C., Shell, B.K., Papadopoulos, N., Vogelstein, B., and Kinzler, K.W. 1995. Genomic organization of the human PMS2 gene family. *Genomics* **30**:195-206.

10 20. Nicolaides, N.C., Palombo, F., Kinzler, K.W., Vogelstein, B., and Jiricny, J. 1996. Molecular cloning of the N-terminus of GTBP. *Genomics* **31**:395-397.

15 21. Palombo, F., Hughes, M., Jiricny, J., Truong, O., Hsuan, J. 1994. Mismatch repair and cancer. *Nature* **36**:417.

20 22. Palombo, F., Gallinari, P., Iaccarino, I., Lettieri, T., Hughes, M.A., Truong, O., Hsuan, J.J., and Jiricny, J. 1995. GTBP, a 160-kilodalton protein essential for mismatch-binding activity in human cells. *Science* **268**:1912-1914.

25 23. Pang, Q., Prolla, T.A., and Liskay, R.M. 1997. Functional domains of the *Saccharomyces cerevisiae* Mlh1p and Pms1p DNA mismatch repair proteins and their relevance to human hereditary nonpolyposis colorectal cancer-associated mutations. *Mol. Cell. Biol.* **17**:4465-4473.

30 24. Papadopoulos, N., Nicolaides, N.C., Wei, Y.F., Carter, K.C., Ruben, S.M., Rosen, C.A., Haseltine, W.A., Fleischmann, R.D., Fraser, C.M., Adams, M.D., Venter, C.J., Dunlop, M.G., Hamilton, S.R., Petersen, G.M., de la Chapelle, A., Vogelstein, B., and kinzler, K.W. 1994. Mutation of a

mutL homolog is associated with hereditary colon cancer. *Science*
263:1625-1629.

25. Parsons, R., Li, G.M., Longley, M.J., Fang, W.H., Papadopolous, N.,
5 Jen, J., de la Chapelle, A., Kinzler, K.W., Vogelstein, B., and Modrich, P.
1993. Hypermutability and mismatch repair deficiency in RER⁺ tumor
cells. *Cell* **75**:1227-1236.
26. Parsons, R., Li, G.M., Longley, M., Modrich, P., Liu, B., Berk, T.,
10 Hamilton, S.R., Kinzler, K.W., and Vogelstein, B. 1995. Mismatch repair
deficiency in phenotypically normal human cells. *Science* **268**:738-740.
27. Perucho, M. 1996. Cancer of the microsatellite mutator phenotype.
Biol Chem. **377**:675-684.
- 15 28. Prolla, T.A, Pang, Q., Alani, E., Kolodner, R.A., and Liskay, R.M.
1994. MLH1, PMS1, and MSH2 Interaction during the initiation of DNA
mismatch repair in yeast. *Science* **264**:1091-1093.
- 20 29. Strand, M., Prolla, T.A., Liskay, R.M., and Petes, T.D. Destabilization
of tracts of simple repetitive DNA in yeast by mutations affecting DNA
mismatch repair. 1993. *Nature* **365**:274-276.
30. Studier, F.W. Use of bacteriophage T7 lysozyme to improve an
25 inducible T7 expression system. **1991**. *J. Mol. Biol.* 219(1):37-44.
31. Su, S.S., Lahue, R.S., Au, K.G., and Modrich, P. 1988. Mismatch
specificity of methyl directed DNA mismatch corrections in vitro. *J. Biol.*
Chem. **263**:6829-6835.

32. Nicolaides NC, Littman SJ, Modrich P, Kinzler KW, Vogelstein B 1998. A naturally occurring hPMS2 mutation can confer a dominant negative mutator phenotype. *Mol Cell Biol* **18**:1635-1641.
- 5 33. Aronshtam A, Marinus MG. 1996. Dominant negative mutator mutations in the mutL gene of *Escherichia coli*. *Nucleic Acids Res* **24**:2498-2504.
- 10 34. Wu TH, Marinus MG. 1994. Dominant negative mutator mutations in the mutS gene of *Escherichia coli*. *J Bacteriol* **176**:5393-400.
- 15 35. Brosh RM Jr, Matson SW. 1995. Mutations in motif II of *Escherichia coli* DNA helicase II render the enzyme nonfunctional in both mismatch repair and excision repair with differential effects on the unwinding reaction. *J Bacteriol* **177**:5612-5621.
36. Studamire B, Quach T, Alani, E. 1998. *Saccharomyces cerevisiae* Msh2p and Msh6p ATPase activities are both required during mismatch repair. *Mol Cell Biol* **18**:7590-7601.
- 20 37. Alani E, Sokolsky T, Studamire B, Miret JJ, Lahue RS. 1997. Genetic and biochemical analysis of Msh2p-Msh6p: role of ATP hydrolysis and Msh2p-Msh6p subunit interactions in mismatch base pair recognition. *Mol Cell Biol* **17**:2436-2447.
- 25 38. Lipkin SM, Wang V, Jacoby R, Banerjee-Basu S, Baxevanis AD, Lynch HT, Elliott RM, and Collins FS. 2000. MLH3: a DNA mismatch repair gene associated with mammalian microsatellite instability. *Nat Genet* **24**:27-35.
- 30 39. Lee CC, Lin HK, Lin JK. 1994. A reverse mutagenicity assay for alkylating agents based on a point mutation in the beta-lactamase gene at the active site serine codon. *Mutagenesis* **9**:401-405.
- 35 40. Vidal A, Abril N, Pueyo C. 1995. DNA repair by Ogt alkyltransferase influences EMS mutational specificity. *Carcinogenesis* **16**:817-821.
41. Fu, K.P., Grace, M.E., Hsiao, C.L., and Hung, P.P. 1988. Elimination of antibiotic-resistant plasmids by quinolone antibiotics. *Chem. Abstracts* **34**(5):415-418.
- 40 42. BiWang, H., ZhiPeng, H., and Xiong, G. 1999. Transformation of *Escherichia coli* and *Bacillus thuringiensis* and their plasmid curing by

electroporation. J. of Fujian Agricultural University 28(1):43-46.

- 5 43. Brosius, J. 1988. Expression vectors employing lambda-, trp-, lac-, and lpp-derived promoters. Biotechnology 10:205-225.